



STIC Search Report

Biotech-Chem Library

STIC Database Tracking Number: 195103

TO: CECILIA JAISLE
Location: REM/4E78/5C18
Art Unit: 1624
Wednesday, July 19, 2006
Case Serial Number: 10/533697

From: John DiNatale
Location: Biotech-Chem Library
REM-1B65
Phone: (571)272-2557

john.dinatale@uspto.gov

Search Notes

Examiner JAISLE,

See attached results.

If you have any questions about this search, or need assistance understanding the MARPAT indexing, feel free to contact me at any time.

Thank you for using STIC search services!

John DiNatale
Technical Information Specialist
STIC Biotech/Chem Library
(571)272-2557

THIS PAGE BLANK (USPTO)

Search history

Jaisle 10/533697

07/19/2006

=> d his full

(FILE 'HOME' ENTERED AT 10:16:46 ON 19 JUL 2006)

FILE 'REGISTRY' ENTERED AT 10:16:52 ON 19 JUL 2006

L1 STRUCTURE UPLOADED
L2 2 SEA SSS SAM L1
D SCA

FILE 'STNGUIDE' ENTERED AT 10:18:18 ON 19 JUL 2006

FILE 'STNGUIDE' ENTERED AT 10:21:32 ON 19 JUL 2006

FILE 'REGISTRY' ENTERED AT 10:28:53 ON 19 JUL 2006

L3 STRUCTURE UPLOADED
L4 2 SEA SSS SAM L3
D SCA

FILE 'STNGUIDE' ENTERED AT 10:31:21 ON 19 JUL 2006

FILE 'REGISTRY' ENTERED AT 10:31:53 ON 19 JUL 2006

L5 STRUCTURE UPLOADED
L6 2 SEA SSS SAM L5
D SCA
L7 16 SEA SSS FUL L5
SAVE TEMP L7 JAI697STRC/A
D STAT QUE L6
D STAT QUE L7
D SCA L7

FILE 'CAPLUS' ENTERED AT 10:36:35 ON 19 JUL 2006

L8 1 SEA ABB=ON PLU=ON L7

FILE 'BEILSTEIN' ENTERED AT 10:37:03 ON 19 JUL 2006

L9 0 SEA SSS SAM L5
L10 0 SEA SSS FUL L5

FILE 'MARPAT' ENTERED AT 10:37:35 ON 19 JUL 2006

FILE 'CAPLUS' ENTERED AT 10:38:10 ON 19 JUL 2006

E US2003-533697 /APPS
E US2005-533697 /APPS
L11 1 SEA ABB=ON PLU=ON US2005-533697 /AP
D SCA
SEL RN

FILE 'REGISTRY' ENTERED AT 10:39:21 ON 19 JUL 2006

L12 30 SEA ABB=ON PLU=ON (110-91-8/BI OR 16182-04-0/BI OR 5028-13-7/
BI OR 6298-19-7/BI OR 663918-38-5/BI OR 684648-96-2/BI OR
684648-97-3/BI OR 684648-98-4/BI OR 684648-99-5/BI OR 684649-00
-1/BI OR 684649-01-2/BI OR 684649-02-3/BI OR 684649-03-4/BI OR
684649-04-5/BI OR 684649-05-6/BI OR 684649-06-7/BI OR 684649-07
-8/BI OR 684649-08-9/BI OR 684649-09-0/BI OR 684649-10-3/BI OR
684649-11-4/BI OR 684649-12-5/BI OR 684649-13-6/BI OR 684649-14
-7/BI OR 684649-15-8/BI OR 684649-16-9/BI OR 684649-17-0/BI OR
684649-18-1/BI OR 684649-19-2/BI OR 684649-20-5/BI)
L13 14 SEA ABB=ON PLU=ON L12 NOT L7
D SCA

FILE 'STNGUIDE' ENTERED AT 10:41:13 ON 19 JUL 2006

FILE 'REGISTRY' ENTERED AT 10:44:06 ON 19 JUL 2006

L14 STRUCTURE UPLOADED
L15 3 SEA SSS SAM L14
D SCA
L16 29 SEA SSS FUL L14
SAVE TEMP L16 JAI697STRD/A
L17 13 SEA ABB=ON PLU=ON L16 NOT L7
D SCA

FILE 'CAPLUS' ENTERED AT 10:48:42 ON 19 JUL 2006

L18 3 SEA ABB=ON PLU=ON L17
L19 0 SEA ABB=ON PLU=ON L11 AND L18

FILE 'BEILSTEIN' ENTERED AT 10:49:29 ON 19 JUL 2006

L20 0 SEA SSS SAM L14
L21 8 SEA SSS FUL L14
L22 1 SEA ABB=ON PLU=ON L21 NOT L18

FILE 'REGISTRY' ENTERED AT 10:51:15 ON 19 JUL 2006

L23 ANALYZE PLU=ON L16 1- LC : 5 TERMS
D
L24 2 SEA ABB=ON PLU=ON L16 NOT CAPLUS/LC
D SCA
L25 7 SEA ABB=ON PLU=ON L16 AND BEILSTEIN/LC
L26 0 SEA ABB=ON PLU=ON L24 AND L25
D LC L24 1-2
D SCA L24
D ED L24 1-2
D COST

FILE 'MARPAT' ENTERED AT 10:55:02 ON 19 JUL 2006

L27 1 SEA SSS SAM L14
L28 49 SEA SSS FUL L14
SAVE TEMP L28 JAI697MARPD/A
L29 1 SEA SUB=L28 SSS SAM L5
L30 24 SEA SUB=L28 SSS FUL L5

FILE 'CAPLUS' ENTERED AT 11:01:21 ON 19 JUL 2006

L31 24 SEA ABB=ON PLU=ON L30
L32 1 SEA ABB=ON PLU=ON L18 AND L31
L33 0 SEA ABB=ON PLU=ON L18 AND L8

FILE 'WPIX' ENTERED AT 11:02:47 ON 19 JUL 2006

L34 1 SEA SSS SAM L14
D SCA
L35 16 SEA SSS FUL L14
D SCA

FILE 'STNGUIDE' ENTERED AT 11:04:04 ON 19 JUL 2006

FILE 'WPIX' ENTERED AT 11:04:29 ON 19 JUL 2006

L36 2 SEA ABB=ON PLU=ON L35/DCR
SEL SDCN L35
EDIT E31-E46 /SDCN /DCN
L37 2 SEA ABB=ON PLU=ON (RAEA00/DCN OR RAEA01/DCN OR RAEA02/DCN OR
RAEA03/DCN OR RAEA04/DCN OR RAEA05/DCN OR RAEA07/DCN OR
RAEA08/DCN OR RAE9ZS/DCN OR RAE9ZT/DCN OR RAE9ZU/DCN OR
RAE9ZV/DCN OR RAE9ZW/DCN OR RAE9ZX/DCN OR RAE9ZY/DCN OR
RAFSCY/DCN)

SEL DCSE L35
 EDIT E47-E62 /DCSE /DCRE
 L38 2 SEA ABB=ON PLU=ON (902542-0-0-0/DCRE OR 902543-0-0-0/DCRE OR
 902544-1-0-0/DCRE OR 902544-2-0-0/DCRE OR 902546-0-0-0/DCRE OR
 902547-0-0-0/DCRE OR 902548-1-0-0/DCRE OR 902548-2-0-0/DCRE OR
 902550-1-0-0/DCRE OR 902551-0-0-0/DCRE OR 902552-1-0-0/DCRE OR
 902552-2-0-0/DCRE OR 902553-0-0-0/DCRE OR 902557-0-0-0/DCRE OR
 902558-0-0-0/DCRE OR 976436-1-0-0/DCRE)
 L39 2 SEA ABB=ON PLU=ON (L36 OR L37 OR L38)

FILE 'STNGUIDE' ENTERED AT 11:07:27 ON 19 JUL 2006

FILE 'CAPLUS' ENTERED AT 11:07:54 ON 19 JUL 2006
 L40 4 SEA ABB=ON PLU=ON L16

FILE 'BABS' ENTERED AT 11:08:44 ON 19 JUL 2006

FILE 'BEILSTEIN' ENTERED AT 11:09:10 ON 19 JUL 2006
 SEL BABSAN L21

FILE 'BABS' ENTERED AT 11:09:30 ON 19 JUL 2006
 L41 1 SEA ABB=ON PLU=ON 5564377/BABSAN

FILE 'CAPLUS' ENTERED AT 11:10:41 ON 19 JUL 2006

FILE 'HCAPLUS' ENTERED AT 11:11:51 ON 19 JUL 2006
 L42 4 SEA ABB=ON PLU=ON L16
 L43 275 SEA ABB=ON PLU=ON RAULT S?/AU
 L44 256 SEA ABB=ON PLU=ON LANCELOT J?/AU
 L45 212 SEA ABB=ON PLU=ON KOPP M?/AU
 L46 121 SEA ABB=ON PLU=ON CAIGNARD D?/AU
 L47 516 SEA ABB=ON PLU=ON PFEIFFER B?/AU
 L*** DEL 107 S RENARD P//AU
 L48 511 SEA ABB=ON PLU=ON RENARD P?/AU
 L49 2 SEA ABB=ON PLU=ON L43 AND L44 AND L45 AND L46 AND L47 AND
 L48
 L50 86 SEA ABB=ON PLU=ON L43 AND (L44 OR L45 OR L46 OR L47 OR L48)
 L51 14 SEA ABB=ON PLU=ON L44 AND (L45 OR L46 OR L47 OR L48)
 L52 4 SEA ABB=ON PLU=ON L45 AND (L46 OR L47 OR L48)
 L53 91 SEA ABB=ON PLU=ON L46 AND (L47 OR L48)
 L54 128 SEA ABB=ON PLU=ON L47 AND L48
 L55 269 SEA ABB=ON PLU=ON (L50 OR L51 OR L52 OR L53 OR L54)
 L56 21 SEA ABB=ON PLU=ON L50 AND (L51 OR L52 OR L53 OR L54)
 L57 10 SEA ABB=ON PLU=ON L51 AND (L52 OR L53 OR L54)
 L58 4 SEA ABB=ON PLU=ON L52 AND (L53 OR L54)
 L59 21 SEA ABB=ON PLU=ON (L56 OR L57 OR L58)

FILE 'MARPAT' ENTERED AT 11:16:14 ON 19 JUL 2006

FILE 'HCAPLUS' ENTERED AT 11:16:28 ON 19 JUL 2006
 L60 1 SEA ABB=ON PLU=ON L59 AND L42

FILE 'MARPAT' ENTERED AT 11:16:43 ON 19 JUL 2006

FILE 'HCAPLUS' ENTERED AT 11:16:54 ON 19 JUL 2006
 L61 1 SEA ABB=ON PLU=ON (L43 OR L44 OR L45 OR L46 OR L47 OR L48)
 AND L42

FILE 'MARPAT' ENTERED AT 11:17:33 ON 19 JUL 2006
 L62 21 SEA ABB=ON PLU=ON L43

L63 14 SEA ABB=ON PLU=ON L44
L64 4 SEA ABB=ON PLU=ON L45
L65 55 SEA ABB=ON PLU=ON L46
L66 41 SEA ABB=ON PLU=ON L47
L67 134 SEA ABB=ON PLU=ON L48
L68 1 SEA ABB=ON PLU=ON (L62 OR L63 OR L64 OR L65 OR L66 OR L67)
AND L30

FILE 'WPIX' ENTERED AT 11:18:26 ON 19 JUL 2006

L69 26 SEA ABB=ON PLU=ON RAULT S?/AU
L70 22 SEA ABB=ON PLU=ON LANCELOT J?/AU
L71 54 SEA ABB=ON PLU=ON KOPP M?/AU
L72 58 SEA ABB=ON PLU=ON CAIGNARD D?/AU
L73 128 SEA ABB=ON PLU=ON PFEIFFER B?/AU
L74 239 SEA ABB=ON PLU=ON RENARD P?/AU
L75 1 SEA ABB=ON PLU=ON L39 AND (L69 OR L70 OR L71 OR L72 OR L73
OR L74)
L76 91 SEA ABB=ON PLU=ON (L50 OR L51 OR L52 OR L53 OR L54)
L77 10 SEA ABB=ON PLU=ON (L56 OR L57 OR L58)

FILE 'STNGUIDE' ENTERED AT 11:20:11 ON 19 JUL 2006

FILE 'WPIX' ENTERED AT 11:20:34 ON 19 JUL 2006

L78 1 SEA ABB=ON PLU=ON L77 AND L39

FILE 'STNGUIDE' ENTERED AT 11:20:46 ON 19 JUL 2006

FILE 'REGISTRY' ENTERED AT 11:22:09 ON 19 JUL 2006
D STAT QUE L16

FILE 'HCAPLUS' ENTERED AT 11:22:23 ON 19 JUL 2006

D QUE NOS L59

D QUE NOS L61

L79 21 SEA ABB=ON PLU=ON L59 OR L61

FILE 'MARPAT' ENTERED AT 11:23:27 ON 19 JUL 2006

D QUE NOS L68

FILE 'WPIX' ENTERED AT 11:23:55 ON 19 JUL 2006

D QUE NOS L78

D QUE NOS L77

D QUE NOS L75

L80 10 SEA ABB=ON PLU=ON L78 OR L77 OR L75

FILE 'STNGUIDE' ENTERED AT 11:25:19 ON 19 JUL 2006

FILE 'HCAPLUS, WPIX, MARPAT' ENTERED AT 11:25:45 ON 19 JUL 2006

L81 23 DUP REM L79 L80 L68 (9 DUPLICATES REMOVED)

ANSWERS '1-21' FROM FILE HCAPLUS

ANSWERS '22-23' FROM FILE WPIX

D IBIB ABS HITSTR L81 1-23

FILE 'STNGUIDE' ENTERED AT 11:27:06 ON 19 JUL 2006

FILE 'REGISTRY' ENTERED AT 11:27:26 ON 19 JUL 2006

D STAT QUE L42

L82 29 SEA ABB=ON PLU=ON L42 NOT L79

FILE 'STNGUIDE' ENTERED AT 11:28:34 ON 19 JUL 2006

D COST

D COST FULL

FILE 'STNGUIDE' ENTERED AT 11:30:59 ON 19 JUL 2006

FILE 'HCAPLUS' ENTERED AT 11:31:17 ON 19 JUL 2006

D QUE NOS L42

L83 3 SEA ABB=ON PLU=ON L42 NOT L79

FILE 'BABS' ENTERED AT 11:32:30 ON 19 JUL 2006

D STAT QUE L41

FILE 'BEILSTEIN' ENTERED AT 11:32:55 ON 19 JUL 2006

D STAT QUE L22

FILE 'WPIX' ENTERED AT 11:33:26 ON 19 JUL 2006

D STAT QUE L39

L84 1 SEA ABB=ON PLU=ON L39 NOT L80

FILE 'MARPAT' ENTERED AT 11:33:55 ON 19 JUL 2006

D STAT QUE L30

L85 23 SEA ABB=ON PLU=ON L30 NOT L68

FILE 'STNGUIDE' ENTERED AT 11:34:23 ON 19 JUL 2006

FILE 'HCAPLUS, WPIX, BABS, MARPAT, BEILSTEIN' ENTERED AT 11:35:09 ON 19 JUL 2006

L86 26 DUP REM L83 L84 L41 L85 L22 (3 DUPLICATES REMOVED)

ANSWERS '1-3' FROM FILE HCAPLUS

ANSWERS '4-25' FROM FILE MARPAT

ANSWER '26' FROM FILE BEILSTEIN

D IBIB ABS HITSTR L86 1-3

D IBIB ABS HIT L86 4-25

D IDE ALLREF L86 26

FILE HOME

FILE REGISTRY

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3

DICTIONARY FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

FILE STNGUIDE

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jul 17, 2006 (20060717/UP).

FILE CAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2006 VOL 145 ISS 4
FILE LAST UPDATED: 18 Jul 2006 (20060718/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

FILE BEILSTEIN
FILE LAST UPDATED ON JUNE 16, 2006

FILE COVERS 1771 TO 2006.

FILE CONTAINS 9,606,495 SUBSTANCES

>>>PLEASE NOTE: Reaction Data and substance data are stored in separate documents and can not be searched together in one query. Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a compounds with available reaction information by combining with PRE/FA, REA/FA or more generally with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be searched as reaction partner BRNs Reactant BRN (RX.RBRN) or Product BRN (RX.PBRN).<<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

* PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
* SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
* ARE BASED ON THE HIGHEST PRICE CATEGORY. THEREFORE; THESE *
* ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
* FOR PRICE INFORMATION SEE HELP COST *

NEW

* PATENT NUMBERS (PN) AND BABS ACCESSION NUMBERS (BABSAN) CAN NOW BE SEARCHED, SELECTED AND TRANSFERRED.
* NEW DISPLAY FORMATS ALLREF, ALLP AND BABSAN SHOW ALL REFERENCES, ALL PATENT REFERENCES, OR ALL BABS ACCESSION NUMBERS FOR A COMPOUND AT A GLANCE.

FILE MARPAT
FILE CONTENT: 1961-PRESENT VOL 145 ISS 1 (20060714/ED)

SOME MARPAT RECORDS ARE DERIVED FROM INPI DATA FOR 1961-1987

MOST RECENT CITATIONS FOR PATENTS FROM MAJOR ISSUING AGENCIES
(COVERAGE TO THESE DATES IS NOT COMPLETE) :

US 2006118302 08 JUN 2006
DE 102004053653 04 MAY 2006
EP 1653548 03 MAY 2006
JP 2006112980 27 APR 2006
WO 2006053912 26 MAY 2006
GB 2419594 03 MAY 2006
FR 2877004 28 APR 2006
RU 2275374 27 APR 2006
CA 2518664 10 MAR 2006

Expanded G-group definition display now available.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

FILE WPIX

FILE LAST UPDATED: 14 JUL 2006 <20060714/UP>
MOST RECENT DERWENT UPDATE: 200645 <200645/DW>
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,
PLEASE VISIT:
http://www.stn-international.de/training_center/patents/stn_guide.pdf <

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE
<http://scientific.thomson.com/support/patents/coverage/latestupdates/>

>>> PLEASE BE AWARE OF THE NEW IPC REFORM IN 2006, SEE
http://www.stn-international.de/stndatabases/details/ipc_reform.html and
<http://scientific.thomson.com/media/scpdf/ipcrdwpi.pdf> <<<

>>> FOR FURTHER DETAILS ON THE FORTHCOMING DERWENT WORLD PATENTS
INDEX ENHANCEMENTS PLEASE VISIT:
http://www.stn-international.de/stndatabases/details/dwpi_r.html <<<

FILE BABS

FILE LAST UPDATED: 15 JUN 2006 <20060615/UP>
FILE COVERS 1980 TO DATE.

FILE HCAPLUS



Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2006 VOL 145 ISS 4
FILE LAST UPDATED: 18 Jul 2006 (20060718/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=>



chain nodes :

11 13 14 15 16 22 25

ring nodes :

1 2 3 4 5 6 7 8 9 10

ring/chain nodes :

12 17

chain bonds :

7-11 9-12 14-15

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 9-10

exact/norm bonds :

5-7 6-10 7-8 7-11 8-9 9-10 9-12 14-15

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6

G1: [*1], [*2], [*3], [*4], [*5]

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom
12:Atom 13:CLASS 14:Atom 15:CLASS 16:CLASS 17:Atom 22:CLASS 25:CLASS 26:CLASS

Generic attributes :



16:

Saturation : Unsaturated

Element Count :

Node 22: Unlimited

THIS PAGE BLANK (USPTO)



chain nodes :

11 13 14 15 16 22 25

ring nodes :

1 2 3 4 5 6 7 8 9 10

ring/chain nodes :

12 17

chain bonds :

7-11 9-12 14-15

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 9-10

exact/norm bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 7-11 8-9 9-10 9-12 14-15

G1: [*1], [*2], [*3], [*4], [*5]

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom
12:Atom 13:CLASS 14:Atom 15:CLASS 16:CLASS 17:Atom 22:CLASS 25:CLASS 28:CLASS

Generic attributes :

16:

Saturation : Unsaturated

Element Count :

Node 22: Unlimited

N,N1

THIS PAGE BLANK (USPTO)


FILE 'REGISTRY' ENTERED AT 11:22:09 ON 19 JUL 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

```
STRUCTURE FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3
DICTIONARY FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3
```

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

```
=> d stat que L16
L14          STR
```

 x^1
$$Ak \dots O \quad 2$$


G1 [@1] , [@2] , [@3] , [@4] , [@5]

Structure attributes must be viewed using STN Express query preparation.
L16 29 SEA FILE=REGISTRY SSS FUL L14

STRUCTURE QUERY

```
100.0% PROCESSED      712 ITERATIONS
SEARCH TIME: 00.00.01
```

29 ANSWERS

=> file hcaplus

FILE 'HCAPLUS' ENTERED AT 11:22:23 ON 19 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

AUTHOR
SEARCH

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2006 VOL 145 ISS 4

FILE LAST UPDATED: 18 Jul 2006 (20060718/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

'OBI' IS DEFAULT SEARCH FIELD FOR 'HCAPLUS' FILE

=> d que nos L59

L43	275	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	RAULT S?/AU
L44	256	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	LANCELOT J?/AU
L45	212	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	KOPP M?/AU
L46	121	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	CAIGNARD D?/AU
L47	516	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	PFEIFFER B?/AU
L48	511	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	RENARD P?/AU
L50	86	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L43 AND (L44 OR L45 OR L46 OR L47 OR L48)
L51	14	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L44 AND (L45 OR L46 OR L47 OR L48)
L52	4	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L45 AND (L46 OR L47 OR L48)
L53	91	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L46 AND (L47 OR L48)
L54	128	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L47 AND L48
L56	21	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L50 AND (L51 OR L52 OR L53 OR L54)
L57	10	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L51 AND (L52 OR L53 OR L54)
L58	4	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L52 AND (L53 OR L54)
L59	21	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	(L56 OR L57 OR L58)

=> d que nos L61

L14		STR				
L16	29	SEA	FILE=REGISTRY	SSS	FUL	L14
L42	4	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L16
L43	275	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	RAULT S?/AU
L44	256	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	LANCELOT J?/AU
L45	212	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	KOPP M?/AU
L46	121	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	CAIGNARD D?/AU
L47	516	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	PFEIFFER B?/AU
L48	511	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	RENARD P?/AU
L61	1	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	(L43 OR L44 OR L45 OR L46 OR

L47 OR L48) AND L42

=> s L59 or L61
L79 21 L59 OR L61

=> file marpat
FILE 'MARPAT' ENTERED AT 11:23:27 ON 19 JUL 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

FILE CONTENT: 1961-PRESENT VOL 145 ISS 1 (20060714/ED)

SOME MARPAT RECORDS ARE DERIVED FROM INPI DATA FOR 1961-1987

MOST RECENT CITATIONS FOR PATENTS FROM MAJOR ISSUING AGENCIES
(COVERAGE TO THESE DATES IS NOT COMPLETE):

US	2006118302	08 JUN 2006
DE	102004053653	04 MAY 2006
EP	1653548	03 MAY 2006
JP	2006112980	27 APR 2006
WO	2006053912	26 MAY 2006
GB	2419594	03 MAY 2006
FR	2877004	28 APR 2006
RU	2275374	27 APR 2006
CA	2518664	10 MAR 2006

Expanded G-group definition display now available.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

=> d que nos L68

L5	STR
L14	STR
L28	49 SEA FILE=MARPAT SSS FUL L14
L30	24 SEA FILE=MARPAT SUB=L28 SSS FUL L5
L43	275 SEA FILE=HCAPLUS ABB=ON PLU=ON RAULT S?/AU
L44	256 SEA FILE=HCAPLUS ABB=ON PLU=ON LANCELOT J?/AU
L45	212 SEA FILE=HCAPLUS ABB=ON PLU=ON KOPP M?/AU
L46	121 SEA FILE=HCAPLUS ABB=ON PLU=ON CAIGNARD D?/AU
L47	516 SEA FILE=HCAPLUS ABB=ON PLU=ON PFEIFFER B?/AU
L48	511 SEA FILE=HCAPLUS ABB=ON PLU=ON RENARD P?/AU
L62	21 SEA FILE=MARPAT ABB=ON PLU=ON L43
L63	14 SEA FILE=MARPAT ABB=ON PLU=ON L44
L64	4 SEA FILE=MARPAT ABB=ON PLU=ON L45
L65	55 SEA FILE=MARPAT ABB=ON PLU=ON L46
L66	41 SEA FILE=MARPAT ABB=ON PLU=ON L47
L67	134 SEA FILE=MARPAT ABB=ON PLU=ON L48
L68	1 SEA FILE=MARPAT ABB=ON PLU=ON (L62 OR L63 OR L64 OR L65 OR L66 OR L67) AND L30

=> file wpix
FILE 'WPIX' ENTERED AT 11:23:55 ON 19 JUL 2006
COPYRIGHT (C) 2006 THE THOMSON CORPORATION

FILE LAST UPDATED: 14 JUL 2006 <20060714/UP>

MOST RECENT DERWENT UPDATE: 200645 <200645/DW>
 DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,
 PLEASE VISIT:
http://www.stn-international.de/training_center/patents/stn_guide.pdf <

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE
<http://scientific.thomson.com/support/patents/coverage/latestupdates/>

>>> PLEASE BE AWARE OF THE NEW IPC REFORM IN 2006, SEE
http://www.stn-international.de/stndatabases/details/ipc_reform.html and
<http://scientific.thomson.com/media/scpdf/ipcrdwpf.pdf> <<<

>>> FOR FURTHER DETAILS ON THE FORTHCOMING DERWENT WORLD PATENTS
 INDEX ENHANCEMENTS PLEASE VISIT:
http://www.stn-international.de/stndatabases/details/dwpi_r.html <<<
 'BIX' IS DEFAULT SEARCH FIELD FOR 'WPIX' FILE

=> d que nos L78

```

L14          STR
L35          16 SEA FILE=WPIX SSS FUL L14
L36          2 SEA FILE=WPIX ABB=ON PLU=ON L35/DCR
L37          2 SEA FILE=WPIX ABB=ON PLU=ON (RAEA00/DCN OR RAEA01/DCN OR
          RAEA02/DCN OR RAEA03/DCN OR RAEA04/DCN OR RAEA05/DCN OR
          RAEA07/DCN OR RAEA08/DCN OR RAE9ZS/DCN OR RAE9ZT/DCN OR
          RAE9ZU/DCN OR RAE9ZV/DCN OR RAE9ZW/DCN OR RAE9ZX/DCN OR
          RAE9ZY/DCN OR RAFSCY/DCN)
L38          2 SEA FILE=WPIX ABB=ON PLU=ON (902542-0-0-0/DCRE OR 902543-0-0-
          0/DCRE OR 902544-1-0-0/DCRE OR 902544-2-0-0/DCRE OR 902546-0-0-
          0/DCRE OR 902547-0-0-0/DCRE OR 902548-1-0-0/DCRE OR 902548-2-0-
          0/DCRE OR 902550-1-0-0/DCRE OR 902551-0-0-0/DCRE OR 902552-1-0-
          0/DCRE OR 902552-2-0-0/DCRE OR 902553-0-0-0/DCRE OR 902557-0-0-
          0/DCRE OR 902558-0-0-0/DCRE OR 976436-1-0-0/DCRE)
L39          2 SEA FILE=WPIX ABB=ON PLU=ON (L36 OR L37 OR L38)
L43          275 SEA FILE=HCAPLUS ABB=ON PLU=ON RAULT S?/AU
L44          256 SEA FILE=HCAPLUS ABB=ON PLU=ON LANCELOT J?/AU
L45          212 SEA FILE=HCAPLUS ABB=ON PLU=ON KOPP M?/AU
L46          121 SEA FILE=HCAPLUS ABB=ON PLU=ON CAIGNARD D?/AU
L47          516 SEA FILE=HCAPLUS ABB=ON PLU=ON PFEIFFER B?/AU
L48          511 SEA FILE=HCAPLUS ABB=ON PLU=ON RENARD P?/AU
L50          86 SEA FILE=HCAPLUS ABB=ON PLU=ON L43 AND (L44 OR L45 OR L46 OR
          L47 OR L48)
L51          14 SEA FILE=HCAPLUS ABB=ON PLU=ON L44 AND (L45 OR L46 OR L47 OR
          L48)
L52          4 SEA FILE=HCAPLUS ABB=ON PLU=ON L45 AND (L46 OR L47 OR L48)
L53          91 SEA FILE=HCAPLUS ABB=ON PLU=ON L46 AND (L47 OR L48)
L54          128 SEA FILE=HCAPLUS ABB=ON PLU=ON L47 AND L48
L56          21 SEA FILE=HCAPLUS ABB=ON PLU=ON L50 AND (L51 OR L52 OR L53 OR
          L54)
L57          10 SEA FILE=HCAPLUS ABB=ON PLU=ON L51 AND (L52 OR L53 OR L54)
L58          4 SEA FILE=HCAPLUS ABB=ON PLU=ON L52 AND (L53 OR L54)
L77          10 SEA FILE=WPIX ABB=ON PLU=ON (L56 OR L57 OR L58)
L78          1 SEA FILE=WPIX ABB=ON PLU=ON L77 AND L39

```

=> d que nos L77

```

L43          275 SEA FILE=HCAPLUS ABB=ON PLU=ON RAULT S?/AU
L44          256 SEA FILE=HCAPLUS ABB=ON PLU=ON LANCELOT J?/AU
L45          212 SEA FILE=HCAPLUS ABB=ON PLU=ON KOPP M?/AU

```

L46 121 SEA FILE=HCAPLUS ABB=ON PLU=ON CAIGNARD D?/AU
 L47 516 SEA FILE=HCAPLUS ABB=ON PLU=ON PFEIFFER B?/AU
 L48 511 SEA FILE=HCAPLUS ABB=ON PLU=ON RENARD P?/AU
 L50 86 SEA FILE=HCAPLUS ABB=ON PLU=ON L43 AND (L44 OR L45 OR L46 OR
 L47 OR L48)
 L51 14 SEA FILE=HCAPLUS ABB=ON PLU=ON L44 AND (L45 OR L46 OR L47 OR
 L48)
 L52 4 SEA FILE=HCAPLUS ABB=ON PLU=ON L45 AND (L46 OR L47 OR L48)
 L53 91 SEA FILE=HCAPLUS ABB=ON PLU=ON L46 AND (L47 OR L48)
 L54 128 SEA FILE=HCAPLUS ABB=ON PLU=ON L47 AND L48
 L56 21 SEA FILE=HCAPLUS ABB=ON PLU=ON L50 AND (L51 OR L52 OR L53 OR
 L54)
 L57 10 SEA FILE=HCAPLUS ABB=ON PLU=ON L51 AND (L52 OR L53 OR L54)
 L58 4 SEA FILE=HCAPLUS ABB=ON PLU=ON L52 AND (L53 OR L54)
 L77 10 SEA FILE=WPIX ABB=ON PLU=ON (L56 OR L57 OR L58)

=> d que nos L75

L14 STR
 L35 16 SEA FILE=WPIX SSS FUL L14
 L36 2 SEA FILE=WPIX ABB=ON PLU=ON L35/DCR
 L37 2 SEA FILE=WPIX ABB=ON PLU=ON (RAEA00/DCN OR RAEA01/DCN OR
 RAEA02/DCN OR RAEA03/DCN OR RAEA04/DCN OR RAEA05/DCN OR
 RAEA07/DCN OR RAEA08/DCN OR RAE9ZS/DCN OR RAE9ZT/DCN OR
 RAE9ZU/DCN OR RAE9ZV/DCN OR RAE9ZW/DCN OR RAE9ZX/DCN OR
 RAE9ZY/DCN OR RAFSCY/DCN)
 L38 2 SEA FILE=WPIX ABB=ON PLU=ON (902542-0-0-0/DCRE OR 902543-0-0-
 0/DCRE OR 902544-1-0-0/DCRE OR 902544-2-0-0/DCRE OR 902546-0-0-
 0/DCRE OR 902547-0-0-0/DCRE OR 902548-1-0-0/DCRE OR 902548-2-0-
 0/DCRE OR 902550-1-0-0/DCRE OR 902551-0-0-0/DCRE OR 902552-1-0-
 0/DCRE OR 902552-2-0-0/DCRE OR 902553-0-0-0/DCRE OR 902557-0-0-
 0/DCRE OR 902558-0-0-0/DCRE OR 976436-1-0-0/DCRE)
 L39 2 SEA FILE=WPIX ABB=ON PLU=ON (L36 OR L37 OR L38)
 L69 26 SEA FILE=WPIX ABB=ON PLU=ON RAULT S?/AU
 L70 22 SEA FILE=WPIX ABB=ON PLU=ON LANCELOT J?/AU
 L71 54 SEA FILE=WPIX ABB=ON PLU=ON KOPP M?/AU
 L72 58 SEA FILE=WPIX ABB=ON PLU=ON CAIGNARD D?/AU
 L73 128 SEA FILE=WPIX ABB=ON PLU=ON PFEIFFER B?/AU
 L74 239 SEA FILE=WPIX ABB=ON PLU=ON RENARD P?/AU
 L75 1 SEA FILE=WPIX ABB=ON PLU=ON L39 AND (L69 OR L70 OR L71 OR
 L72 OR L73 OR L74)

=> s L78 or L77 or L75

L80 10 L78 OR L77 OR L75

=> => dup rem L79 L80 L68

FILE 'HCAPLUS' ENTERED AT 11:25:45 ON 19 JUL 2006
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 11:25:45 ON 19 JUL 2006
 COPYRIGHT (C) 2006 THE THOMSON CORPORATION



FILE 'MARPAT' ENTERED AT 11:25:45 ON 19 JUL 2006
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2006 American Chemical Society (ACS)
 PROCESSING COMPLETED FOR L79

PROCESSING COMPLETED FOR L80
 PROCESSING COMPLETED FOR L68
 L81 23 DUP REM L79 L80 L68 (9 DUPLICATES REMOVED)
 ANSWERS '1-21' FROM FILE HCAPLUS
 ANSWERS '22-23' FROM FILE WPIX

=> d ibib abs hitstr L81 1-23

this case
 L81 ANSWER 1 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 1
 ACCESSION NUMBER: 2004:370797 HCAPLUS
 DOCUMENT NUMBER: 140:375180
 TITLE: Preparation of pyridopyrimidinones as hypolipemic agents
 INVENTOR(S): Rault, Sylvain; Lancelot, Jean
 Charles; Kopp, Marina; Caignard,
 Daniel Henri; Pfeiffer, Bruno;
 Renard, Pierre
 PATENT ASSIGNEE(S): Les Laboratoires Servier, Fr.
 SOURCE: Fr. Demande, 22 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2846657	A1	20040507	FR 2002-13804	20021105
FR 2846657	B1	20041224		
CA 2503995	AA	20040527	CA 2003-2503995	20031104
WO 2004043956	A1	20040527	WO 2003-FR3274	20031104
WO 2004043956	C1	20050609		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003292320	A1	20040603	AU 2003-292320	20031104
EP 1560826	A1	20050810	EP 2003-767886	20031104
EP 1560826	B1	20060412		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003015801	A	20050920	BR 2003-15801	20031104
CN 1711260	A	20051221	CN 2003-80102713	20031104
JP 2006512317	T2	20060413	JP 2004-550726	20031104
AT 323092	E	20060415	AT 2003-767886	20031104
US 2005288311	A1	20051229	US 2005-533697	20050503
NO 2005002683	A	20050603	NO 2005-2683	20050603
PRIORITY APPLN. INFO.:			FR 2002-13804	A 20021105
			WO 2003-FR3274	W 20031104
OTHER SOURCE(S):	MARPAT 140:375180			
GI				



AB Title compds. I [wherein R1, R2 = independently H, alkyl, or R1NR2 = heterocycle; R3 = halo, alkoxy, (un)substituted aryl; R4 = H, NH2 and derivs.; their enantiomers, diastereoisomers, tautomers, and their addition salts with a pharmaceutically acceptable acid or base] were prepared as hypolipemic agents. For example, II was prepared by reacting 3-amino-2-azepano-6-(4-morpholinyl)pyridine with ethoxycarbonyl isothiocyanate in the presence of thiourea in DMF, alkylation with DIPA in the presence of HgCl2 (no isolation of intermediate), and cyclization in DMF at reflux for 2 h. II at 125 mg/kg and metformin at 250 mg/kg reduced triglycerides to the same level in vivo in obese mice using the insulin resistance associated with obesity model. I activated or inhibited certain kinases (no data). Thus, I are useful for treating cancer, diabetes type II, obesity, hyperlipidemia, hypercholesterolemia, cardiovascular complications, arthrosis, and hypertension.


IT **684648-96-2P**, 8-(1-Azocanyl)-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684648-98-4P**, 8-(4-Thiomorpholinyl)-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-00-1P**, 8-[(4 α ,8 α)Octahydro-1(2H)-quinolinyl]-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-02-3P**, 8-[(4 α ,8 α)Octahydro-1(2H)-quinolinyl]-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-03-4P**, 6,8-Di(1-azepanyl)-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-05-6P**, 8-(1-Azepanyl)-2-(dipropylamino)-6-(4-morpholinyl)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-07-8P**, 8-(1-Azepanyl)-2,6-di(4-morpholinyl)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-11-4P**, 2-Amino-8-[(3 α ,5 β)-3,5-dimethylmorpholin-4-yl]pyrido[3,4-d]pyrimidin-4(3H)-one **684649-12-5P**, 2-Amino-8-[(3 α ,5 α)-3,5-dimethylmorpholinyl]pyrido[3,4-d]pyrimidin-4(3H)-one **684649-13-6P**, 8-[(3 α ,5 α)-3,5-Dimethylmorpholinyl]-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-14-7P**, 8-[(3 α ,5 β)-3,5-Dimethylmorpholinyl]-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-15-8P**,

8-[(3 α ,5 α)-3,5-Dimethylmorpholinyl]-2-(4-morpholinyl)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-16-9P**,
 2-Amino-8-(1-azepanyl)-6-(4-morpholinyl)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-17-0P**, 8-Chloro-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-18-1P**, 2-(Dipropylamino)-8-(1-pyrrolidinyl)pyrido[3,4-d]pyrimidin-4(3H)-one **684649-19-2P**,
 8-(3,4-Dimethoxyphenyl)-2-(dipropylamino)pyrido[3,4-d]pyrimidin-4(3H)-one
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of pyridopyrimidinones as hypolipemic agents)


RN 684648-96-2 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-(hexahydro-1(2H)-azocinyl)- (9CI) (CA INDEX NAME)



RN 684648-98-4 HCAPLUS


CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-(4-thiomorpholinyl)- (9CI) (CA INDEX NAME)



RN 684649-00-1 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-[(4aR,8aR)-octahydro-1(2H)-quinolinyl]-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 684649-02-3 HCAPLUS


CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-[(4aR,8aS)-octahydro-1(2H)-quinolinyl]-, rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 684649-03-4 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-6,8-bis(hexahydro-1H-azepin-1-yl)- (9CI) (CA INDEX NAME)




RN 684649-05-6 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-(hexahydro-1H-azepin-1-yl)-6-(4-morpholinyl)- (9CI) (CA INDEX NAME)



RN 684649-07-8 HCAPLUS


CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-(hexahydro-1H-azepin-1-yl)-2,6-di-4-morpholinyl- (9CI) (CA INDEX NAME)



RN 684649-11-4 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-amino-8-[(3R,5R)-3,5-dimethyl-4-morpholinyl]-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 684649-12-5 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-amino-8-[(3R,5S)-3,5-dimethyl-4-morpholinyl]-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 684649-13-6 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-[(3R,5S)-3,5-dimethyl-4-morpholinyl]-2-(dipropylamino)-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 684649-14-7 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-[(3R,5R)-3,5-dimethyl-4-morpholinyl]-2-(dipropylamino)-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.




RN 684649-15-8 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-[(3R,5S)-3,5-dimethyl-4-morpholinyl]-2-(4-morpholinyl)-, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.




RN 684649-16-9 HCAPLUS
 CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-amino-8-(hexahydro-1H-azepin-1-yl)-6-(4-morpholinyl)- (9CI) (CA INDEX NAME)



RN 684649-17-0 HCAPLUS
 CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-chloro-2-(dipropylamino)- (9CI) (CA INDEX NAME)




RN 684649-18-1 HCAPLUS
 CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 2-(dipropylamino)-8-(1-pyrrolidinyl)- (9CI) (CA INDEX NAME)



RN 684649-19-2 HCAPLUS

CN Pyrido[3,4-d]pyrimidin-4(1H)-one, 8-(3,4-dimethoxyphenyl)-2-(dipropylamino)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 2 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 2004:370796 HCAPLUS

DOCUMENT NUMBER: 140:375173

TITLE: Preparation of imidazopyridines as AMPK activators for treating diabetes and hyperlipidemia

INVENTOR(S): Rault, Sylvain; Lancelot, Jean Charles; Kopp, Marina; Caignard, Daniel Henri; Pfeiffer, Bruno; Renard, Pierre; Bizot Espiard, Jean Guy

PATENT ASSIGNEE(S): Les Laboratoires Servier, Fr.

SOURCE: Fr. Demande, 21 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2846656	A1	20040507	FR 2002-13802	20021105
FR 2846656	B1	20041224		
CA 2504008	AA	20040527	CA 2003-2504008	20031104
WO 2004043957	A1	20040527	WO 2003-FR3277	20031104



W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2003292323 A1 20040603 AU 2003-292323 20031104
 EP 1558612 A1 20050803 EP 2003-767889 20031104
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
 BR 2003015800 A 20050920 BR 2003-15800 20031104
 CN 1711261 A 20051221 CN 2003-80102714 20031104
 JP 2006508111 T2 20060309 JP 2004-550729 20031104
 US 2006069117 A1 20060330 US 2005-533699 20050503
 NO 2005002710 A 20050606 NO 2005-2710 20050606
 PRIORITY APPLN. INFO.: FR 2002-13802 A 20021105
 WO 2003-FR3277 W 20031104

OTHER SOURCE(S): MARPAT 140:375173
 GI



AB Title compds. I [wherein R1 = H, halo, polyhalogeno/alkyl, CN, NO2, hydroxycarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, or dialkylaminocarbonyl; R2 = H, alkyl, (un)substituted hetero/aryl; R3 = H, alkyl; X = (CH2)_n; n = 1-6; A = pyridine ring; their enantiomers, diastereoisomers, and their addition salts with a pharmaceutically acceptable acid or base] were prepared as AMP protein kinase (AMPK) activators for treating diabetes and hyperlipidemia. Thus, II (m.p. = 210°) was prepared by reaction of 3-amino-2-cycloheptylaminopyridine with ethoxycarbonyl isothiocyanate in DMF for 3 h, intramol. cyclization in MeOH in the presence of base, and ethoxycarbonyl deprotection in the presence of gaseous HCl and dioxane at reflux for 12 h. II, at 500 μM, activated AMP kinase after 30 min by 312% compared to 178% activation by 5-aminoimidazole-4-carboxamidoriboside in a cellular model. II at 125 mg/kg and metformin at 250 mg/kg reduced triglycerides to the same level in rats. Thus, I are useful for treating hypercholesterolemia, diabetes,


hyperlipidemia, obesity, and cardiovascular complications.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


10 L81 ANSWER 3 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 3
 ACCESSION NUMBER: 2004:370795 HCAPLUS
 DOCUMENT NUMBER: 140:375078
 TITLE: Preparation of pyridines as hypoglycemic agents for treating diabetes and obesity
 INVENTOR(S): Rault, Sylvain; Kopp, Marina; Lancelot, Jean Charles; Lemaitre, Stephane; Pfeiffer, Bruno; Bizot Espiard, Jean Guy; Renard, Pierre
 PATENT ASSIGNEE(S): Les Laboratoires Servier, Fr.
 SOURCE: Fr. Demande, 19 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2846655	A1	20040507	FR 2002-13801	20021105
FR 2846655	B1	20041224		
WO 2004043947	A1	20040527	WO 2003-FR3275	20031104
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR				
AU 2003292321	A1	20040603	AU 2003-292321	20031104
PRIORITY APPLN. INFO.:			FR 2002-13801	A 20021105
			WO 2003-FR3275	W 20031104

OTHER SOURCE(S): MARPAT 140:375078
 GI



I



II

AB Title compds. I [wherein R1, R2 = independently H, cyclo/alkyl, (un)substituted hetero/aryl, or R1NR2 = (un)substituted (non)aromatic heterocycle; R3 = H, alkyl, alkoxy, OH, CN, NH2, etc.; R4, R5 = independently H, alkyl; X = O, S, NH and derivs. with certain compds. absent; their enantiomers, diastereomers, tautomers, and their addition salts

with a pharmaceutically acceptable acid or base, with provisos] were prepared as hypoglycemic agents for treating diabetes and obesity. For example, II (m.p. = 147°) was prepared by N-alkylation of morpholine with 2-chloronicotinonitrile, and cyclization of nitrile with 1,2-propanediamine in the presence of P2S5. II reduced glycemia in non-diabetic Wistar rats, STZ-induced diabetes Wistar rats, and Zucker rats by 16/1.4 %, 21/1.93 %, and 9/1.24 %, resp. when administered orally at a dose of 10 mg/kg. II was non-toxic at a maximum concentration of 1024 mg/kg.

Thus, I are useful for treating diabetes and obesity.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 4 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 4

ACCESSION NUMBER: 2004:370794 HCAPLUS

DOCUMENT NUMBER: 140:375077

TITLE: Preparation of 2,3-dihydro-4(1H)-pyridinones as memory enhancers and analgesics

INVENTOR(S): **Rault, Sylvain**; Leflemme, Nicolas;
Dallemagne, Patrick; Lestage, Pierre; Lockhart, Brian;
Danober, Laurence; **Pfeiffer, Bruno**;
Renard, Pierre

PATENT ASSIGNEE(S): Les Laboratoires Servier, Fr.

SOURCE: Fr. Demande, 24 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1



PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2846654	A1	20040507	FR 2002-13803	20021105
CA 2503993	AA	20040527	CA 2003-2503993	20031104
WO 2004043952	A1	20040527	WO 2003-FR3276	20031104
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003292322	A1	20040603	AU 2003-292322	20031104
EP 1560825	A1	20050810	EP 2003-767888	20031104
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003015996	A	20050927	BR 2003-15996	20031104
CN 1705660	A	20051207	CN 2003-80101823	20031104
JP 2006508110	T2	20060309	JP 2004-550728	20031104
US 2006019995	A1	20060126	US 2005-533784	20050504
NO 2005002598	A	20050530	NO 2005-2598	20050530
PRIORITY APPLN. INFO.:			FR 2002-13803	A 20021105
			WO 2003-FR3276	W 20031104

OTHER SOURCE(S): MARPAT 140:375077

GI


FR 2846328	A1	20040430	FR 2002-13194	20021023
FR 2846328	B1	20041210		
JP 2004143167	A2	20040520	JP 2003-357410	20031017
US 2004087638	A1	20040506	US 2003-689394	20031020
US 6875788	B2	20050405		
BR 2003004634	A	20040601	BR 2003-4634	20031021
CA 2445817	AA	20040423	CA 2003-2445817	20031022
AT 288432	E	20050215	AT 2003-292634	20031022
NZ 529084	A	20050429	NZ 2003-529084	20031022
PT 1413579	T	20050531	PT 2003-292634	20031022
ES 2236672	T3	20050716	ES 2003-3292634	20031022
AU 2003257606	A1	20040513	AU 2003-257606	20031023
ZA 2003008265	A	20040709	ZA 2003-8265	20031023
HK 1063320	A1	20060504	HK 2004-106086	20040813
US 2005143440	A1	20050630	US 2005-69094	20050301
PRIORITY APPLN. INFO.:			FR 2002-13194	A 20021023
			US 2003-689394	A3 20031020
OTHER SOURCE(S):		MARPAT 140:357344		
GI				




AB Title compds. I [wherein R1 = (un)substituted heteroaryl; R2 = (un)substituted cycloalkyl; R3 = H, alkyl; R4, R5 = independently H, halo, polyhalogeno/alkyl, etc.; their enantiomers, diastereomers, tautomers, and their salts of addition with a pharmaceutically acceptable acid or base; with provisos] were prepared as antidiabetic agents. For example, II was prepared by cyclocondensation of cyclohexyl(3-thienyl)acetonitrile (preparation given) with 1,2-propanediamine in the presence of PS5. II was tested for use as a drug for non-insulin dependent diabetes and hyperlipidemia associated with obesity (glycemia reduced 13 - 18% at 10 mg/kg). I are useful for treating diabetes mellitus type II, obesity, diabetes type I, hyperlipidemia, hypercholesterolemia, and cardiovascular complications.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L81 ANSWER 6 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 6
 ACCESSION NUMBER: 2000:790170 HCAPLUS
 DOCUMENT NUMBER: 133:335170
 TITLE: Preparation of 1-aza-2-alkyl-6-arylcycloalkanes useful as memory enhancers
 INVENTOR(S): Rault, Sylvain; Renault, Olivier; Guillon, Jean; Dallemagne, Patrick; Renard, Pierre; Pfeiffer, Bruno; Lestage, Pierre; Lebrun, Marie-Cecile
 PATENT ASSIGNEE(S): Adir et Compagnie, Fr.; Servier Lab.
 SOURCE: Eur. Pat. Appl., 26 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent



I



II



III

AB Title compds. I [wherein R1 = H, aryl/alkyl, acyl, alkoxycarbonyl, arylalkoxycarbonyl, CF₃; R2 = H, alkyl; X = O or NOR₃; R3 = H, (un)substituted alkyl; Ar = hetero/aryl; their enantiomers, diastereomers isomers, and their addition salts with a pharmaceutically acceptable acid] were prepared as memory enhancers and analgesics. For example, II was prepared by Pd-cross coupling of iodide III with 2-thienylboronic acid, and Boc deprotection. II inhibited abdominal cramps induced by PBQ in mice by 48% when administered i.p..

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 5 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 5

ACCESSION NUMBER: 2004:348013 HCAPLUS
 DOCUMENT NUMBER: 140:357344
 TITLE: Preparation of imidazolines as hypoglycemic agents
 INVENTOR(S): Rault, Sylvain; Kopp, Marina;
 Lancelot, Jean-Charles; Lemaitre, Stephane;
 Caignard, Daniel-Henri; Bizot-espiard,
 Jean-Guy; Renard, Pierre
 PATENT ASSIGNEE(S): Les Laboratoires Servier, Fr.
 SOURCE: Eur. Pat. Appl., 13 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1413579	A1	20040428	EP 2003-292634	20031022
EP 1413579	B1	20050202		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK

LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1050530	A1	20001108	EP 2000-401199	20000502
EP 1050530	B1	20031022		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
FR 2793246	A1	20001110	FR 1999-5600	19990503
FR 2793246	B1	20010629		
JP 2000351766	A2	20001219	JP 2000-127382	20000427
JP 3224376	B2	20000427		
CN 1277194	A	20001220	CN 2000-118158	20000430
CN 1145612	B	20040414		
NO 2000002315	A	20001106	NO 2000-2315	20000502
NO 315850	B1	20031103		
NZ 504299	A	20010126	NZ 2000-504299	20000502
US 6323222	B1	20011127	US 2000-561646	20000502
AT 252558	E	20031115	AT 2000-401199	20000502
PT 1050530	T	20040227	PT 2000-401199	20000502
ES 2209777	T3	20040701	ES 2000-401199	20000502
CA 2308783	AA	20001103	CA 2000-2308783	20000503
CA 2308783	C	20041019		
ZA 2000002151	A	20001107	ZA 2000-2151	20000503
BR 2000002076	A	20010123	BR 2000-2076	20000503
AU 763685	B2	20030731	AU 2000-31324	20000503
HK 1032235	A1	20040930	HK 2001-102865	20010423
US 2002042413	A1	20020411	US 2001-964085	20010926
US 6451789	B2	20020917		

PRIORITY APPLN. INFO.:

FR 1999-5600 A 19990503
 US 2000-561646 A3 20000502

OTHER SOURCE(S):
 GI

MARPAT 133:335170



AB The title compds. I [n = 0, 1; R1 = H, arylalkyl, alkyl, acyl, etc.; R2 = alkyl; X = O, Cl, OR3, SR4, NOR5; Ar = aryl, heteroaryl], memory enhancers, were prepared E.g., 2-(3-chlorophenyl)-6-methyl-2,3-dihydro-4(1H)-pyridinone was prepared

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 7 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 7
 ACCESSION NUMBER: 1996:485727 HCAPLUS
 DOCUMENT NUMBER: 125:142700
 TITLE: Tricyclic oxime ethers process for their preparation

and pharmaceutical compositions containing them

INVENTOR(S) : **Rault, Sylvain**; Robba, Max; **Lancelot, Jean-Charles**; Prunier, Herve; **Renard, Pierre**; **Pfeiffer, Bruno**;
Guardiola-Lemaitre, Beatrice; Rettori, Marie-Claire

PATENT ASSIGNEE(S) : Adir Et Compagnie, Fr.

SOURCE: Eur. Pat. Appl., 45 pp.
CODEN: EPXXDW


DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 718299	A1	19960626	EP 1995-402865	19951219
EP 718299	B1	20000405		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
FR 2728571	A1	19960628	FR 1994-15431	19941222
FR 2728571	B1	19970131		
CA 2165618	AA	19960623	CA 1995-2165618	19951219
CA 2165618	C	20010410		
AT 191483	E	20000415	AT 1995-402865	19951219
PT 718299	T	20000731	PT 1995-402865	19951219
ES 2147271	T3	20000901	ES 1995-402865	19951219
FI 9506136	A	19960623	FI 1995-6136	19951220
AU 9540593	A1	19960627	AU 1995-40593	19951220
AU 693615	B2	19980702		
NO 9505215	A	19960624	NO 1995-5215	19951221
ZA 9510901	A	19960624	ZA 1995-10901	19951221
JP 08231554	A2	19960910	JP 1995-333347	19951221
JP 2937837	B2	19990823		
US 5627203	A	19970506	US 1995-576678	19951221
CN 1131155	A	19960918	CN 1995-120144	19951222
CN 1066449	B	20010530		
CN 1261073	A	20000726	CN 1999-120993	19991203
GR 3033507	T3	20000929	GR 2000-401198	20000525
PRIORITY APPLN. INFO.:			FR 1994-15431	A 19941222
OTHER SOURCE(S) :	MARPAT 125:142700			
GI				



AB The present invention concerns compds. I, in which A represents a thieno group, x and y are independently 0-4, R1 is H, alkyl, alkenyl, cycloalkyl, OH, alkoxy, substituted Ph, phenylalkyl, substituted phenoxy, R2 and R3 are H, alkyl, alkenyl, cycloalkyl, substituted indanyl, substituted Ph, phenylalkyl, or R2 and R3 form azacycloalkyl rings, and their oxalates or fumarates. I, e.g. II (X = NOCHPhCH2CH2NMe2) are prepared from the ketone, e.g. II (X = O), via hydroxyimination followed by O-alkylation, e.g. with

PhCHClCH₂CH₂NMe₂·HCl. I were tested as serotoninergetic receptor antagonists (IC₅₀ 1.1 x 10⁻¹⁰ to 10⁻⁴ M), anxiolytics and antidepressants.

46 L81 ANSWER 8 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 8

ACCESSION NUMBER: 1994:217734 HCAPLUS

DOCUMENT NUMBER: 120:217734

TITLE: Pyrrolothienopyrazine serotonin 5-HT₃ receptor antagonists

INVENTOR(S): Rault, Sylvain; Lancelot, Jean
Charles; Pilo Vincente, Juan Carlos; Robba, Max;
Guardiola-Lemaitre, Beatrice; Renard, Pierre
; Adam, Gerard

PATENT ASSIGNEE(S): ADIR et Compagnie, Fr.

SOURCE: Eur. Pat. Appl., 35 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 573360	A1	19931208	EP 1993-401416	19930603
EP 573360	B1	19980826		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
FR 2691967	A1	19931210	FR 1992-6800	19920605
FR 2691967	B1	19950609		
AT 170187	E	19980915	AT 1993-401416	19930603
ES 2123038	T3	19990101	ES 1993-401416	19930603
CA 2097779	AA	19931206	CA 1993-2097779	19930604
AU 9340059	A1	19931209	AU 1993-40059	19930604
AU 659738	B2	19950525		
ZA 9303942	A	19931230	ZA 1993-3942	19930604
JP 06172363	A2	19940621	JP 1993-134922	19930604
JP 07094460	B4	19951011		

PRIORITY APPLN. INFO.:

FR 1992-6800 A 19920605

OTHER SOURCE(S): MARPAT 120:217734


GI



AB The title compds. I [A = (un)substituted Q₁, (un)substituted Q₂; R₁R₂ form

a double bond and R3 represents a Cl atom, a substituted amine or heterocyclyl group, or no group or R1 may represent a H and R2R3 a :O], which are highly specific serotonin 5-HT3 receptor antagonists (no data), useful in the treatment of depression (no data), stress (no data) psychoses (no data), migraine headache (no data), etc., are prepared and I-containing formulations presented. Thus, 2-(pyrrol-1-yl)-3-thenoyl nitride was refluxed in 1,2-dichlorobenzene, the intermediate heated in the presence of POCl₂, and condensed with 1-methylpiperazine, producing pyrazine II, m.p. 82°.

76
L81 ANSWER 9 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2004:115127 HCAPLUS
DOCUMENT NUMBER: 140:321259
TITLE: Design, synthesis, and evaluation of novel
thienopyrrolizinones as antitubulin agents
AUTHOR(S): Lisowski, Vincent; Leonce, Stephane; Kraus-Berthier,
Laurence; Santos, Jana Sopkova-de Oliveira; Pierre,
Alain; Atassi, Ghanem; **Caignard, Daniel-Henri**
; **Renard, Pierre; Rault, Sylvain**
CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de
Normandie UFR des Sciences Pharmaceutiques, Universite
de Caen, Caen, 14032, Fr.
SOURCE: Journal of Medicinal Chemistry (2004), 47(6),
1448-1464
CODEN: JMCMAR; ISSN: 0022-2623
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 140:321259
GI




AB The structure-activity relationship study of a new 3-aryl-8-H-thieno[2,3-b]pyrrolizin-8-one series of antitubulin agents was described. The pharmacol. results from the National Cancer Institute in vitro human disease oriented tumor cell line screening allowed the identification of 3-(4-methoxyphenyl)-8H-thieno[2,3-b]pyrrolizin-8-one (NSC 676693) as a very efficient antitumor drug in all cancer cell lines tested. This required a definition of structural requirements essential for this antiproliferative activity. Among all analogs synthesized in this study, 3-(3-hydroxy-4-methoxyphenyl)-8H-thieno[2,3-b]pyrrolizin-8-one (I) was the most promising, being 10-fold more potent than NSC 676693. Its activity over a panel of nine tumoral cell lines was in the nanomolar range for all of the histol. types tested, and surprisingly, the resistant KB-A1 cell

line was also sensitive to this compound. Moreover, a flow cytometric study showed that L1210 cells treated by the most potent compds. were arrested in the G2/M phases of the cell cycle with a significant percentage of cells having reinitiated a cycle of DNA synthesis without cell division. This interesting pharmacol. profile, resulting from inhibition of tubulin polymerization, encouraged us to perform preliminary in vivo studies that led to


a new prodrug chemical approach.

REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

460 L81 ANSWER 10 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2002:928395 HCAPLUS
 DOCUMENT NUMBER: 138:170063
 TITLE: New Benzo[h][1,6]naphthyridine and Azepino[3,2-c]quinoline Derivatives as Selective Antagonists of 5-HT₄ Receptors: Binding Profile and Pharmacological Characterization
 AUTHOR(S): Hinschberger, Antoine; Butt, Sabrina; Lelong, Veronique; Boulouard, Michel; Dumuis, Aline; Dauphin, Francois; Bureau, Ronan; Pfeiffer, Bruno; Renard, Pierre; Rault, Sylvain
 CORPORATE SOURCE: ATBI, Universite de Caen, Caen, 14032, Fr.
 SOURCE: Journal of Medicinal Chemistry (2003), 46(1), 138-147
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 138:170063
 GI



I



II

AB Benzo[h][1,6]naphthyridinyloxymethylpiperidines such as I (n = 1),

azepino[3,2-c]quinolinyloxymethylpiperidines such as I (n = 2), and benzonaphthyridines II (R = EtCH₂, Bu) were prepared as potential 5-HT₄ receptor antagonists. Benzonaphthyridinyloxymethylpiperidines and azepinoquinolinyloxymethylpiperidines were prepared by substitution reactions of chlorobenzonaphthyridines and chloroazepinoquinolines with N-substituted 4-piperidinemethanols. The binding of benzonaphthyridinyloxymethylpiperidines and azepinoquinolinyloxymethylpiperidines to the 5-HT₄ receptor decreased markedly when chlorine atom substituents were present on the aromatic ring; N-Pr and N-Bu substituents on the piperidine moiety yield compds. such as I (n = 1) which bound to 5-HT₄ receptors with nanomolar affinities. I (n = 1) acted as an antagonist/low partial agonist of the 5-HT₄ receptor. I (n = 1) also showed potent analgesic activity in mice at doses of 0.01-1 mg/kg.

REFERENCE COUNT: 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


1/b L81 ANSWER 11 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2002:805388 HCAPLUS
DOCUMENT NUMBER: 138:238138
TITLE: A versatile synthesis of 2-amino-4H-pyrido[1,2-a][1,3,5]triazin-4-ones from 2-aminopyridines
AUTHOR(S): Kopp, Marina; Lancelot, Jean-Charles
; Dagdag, Said; Miel, Hugues; Rault, Sylvain
CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de Normandie, Universite de Caen-Basse Normandie, Caen, 14032, Fr.
SOURCE: Journal of Heterocyclic Chemistry (2002), 39(5), 1061-1064
CODEN: JHTCAD; ISSN: 0022-152X
PUBLISHER: HeteroCorporation
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 138:238138
AB The quasi-one pot synthesis of new 2-amino-4H-pyrido[1,2-a]-1,3,5-triazin-4-ones starting from 2-aminopyridine and 2-aminopicolines is herein described in order to obtain a library of cyclic guanidines.
REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

1/b L81 ANSWER 12 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2002:938490 HCAPLUS
DOCUMENT NUMBER: 139:30285
TITLE: Synthesis of New Aromatic Pyrrolo[2,1-c][1,4]benzodiazepines and Pyrrolo[1,2-a]thieno[3,2-e][1,4]diazepines as Anti-tumoral Agents
AUTHOR(S): Lisowski, Vincent; Fabis, Frederic; Pierre, Alain; Caignard, Daniel-Henri; Renard, Pierre
; Rault, Sylvain
CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de Normandie-UFR des Sciences Pharmaceutiques 5, Caen, 14032, Fr.
SOURCE: Journal of Enzyme Inhibition and Medicinal Chemistry (2002), 17(6), 403-407
CODEN: JEIMAZ; ISSN: 1475-6366
PUBLISHER: Taylor & Francis Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 139:30285
AB Diazepine analogs of thieno[2,3-b]pyrrolizin-8-ones were synthesized by aromatization of 2-hydroxypyrrolo[1,2-a]thieno[3,2-e][1,4]diazepines.

These compds. were evaluated in vitro for their antiproliferative activity against the L1210 leukemia cell line. The activity of these compds. was in the micromolar range, the best result being for the mixture of the isomers which showed a 0.35 μM IC50 against cell growth.

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

81 ANSWER 13 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2001:612043 HCAPLUS
 DOCUMENT NUMBER: 135:371658
 TITLE: Design, synthesis and antiproliferative activity of tripentones: A new series of antitubulin agents
 AUTHOR(S): Lisowski, V.; Enguehard, C.; Lancelot, J.-C.; Caignard, D.-H.; Lambel, S.; Leonce, S.; Pierre, A.; Atassi, G.; Renard, P.; Rault, S.
 CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de Normandie, UFR des Sciences Pharmaceutiques, Caen, 14032, Fr.
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2001), 11(16), 2205-2208
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:371658
 GI




I

AB Structure-activity relationship studies of a new series of tripentones (thieno[2,3-b]pyrrolizin-8-ones), led to the prepare of several derivs. with antiproliferative activities. The most promising 3-(3-hydroxy-4-methoxyphenyl)thieno[2,3-b]pyrrolizin-8-one (I) (leukemia L1210, IC50=15 nM) was shown to be a potent inhibitor of tubulin polymerization

REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

81 ANSWER 14 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 2001:779581 HCAPLUS
 DOCUMENT NUMBER: 136:183790
 TITLE: Synthesis of novel pyrazolopyrrolopyrazines, potential analogs of sildenafil
 AUTHOR(S): Kopp, Marina; Lancelot, Jean-Charles; Dallemagne, Patrick; Rault, Sylvain
 CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de Normandie, Caen, 14032, Fr.
 SOURCE: Journal of Heterocyclic Chemistry (2001), 38(5), 1045-1050

PUBLISHER: HeteroCorporation
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 136:183790
GI



AB The pyrazolopyrrolopyrazines I [R = Me, Ph, 4-MeOC6H4, R1 = H; R = Me, R1 = Pr] were prepared as analogs of sildenafil. I were 10-fold less active than sildenafil as PDE-5 inhibitors.

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 15 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2001:177391 HCAPLUS

DOCUMENT NUMBER: 135:28643

TITLE: First tricyclic oximino derivatives as 5-HT3 ligands

AUTHOR(S): Baglin, I.; Daveu, C.; **Lancelot, J. C.**;

Bureau, R.; Dauphin, F.; **Pfeiffer, B.**;

Renard, P.; Delagrang, P.; **Rault, S.**

CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de Normandie, Universite de Caen, Caen, 14032, Fr.

SOURCE: Bioorganic & Medicinal Chemistry Letters (2001), 11(4), 453-457

CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The design and synthesis of a new type of 5-HT3 ligand with subnanomolar affinity are described. The O-dialkylaminoethyloximinothienopyrrolizine structure was deduced from mol. modeling studies by replacement of an amidine moiety by an oximino one.

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L81 ANSWER 16 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:106891 HCAPLUS

DOCUMENT NUMBER: 132:151677

TITLE: Preparation of 8H-thieno-[2,3-b]pyrrolizin-8-ones as anticancer agents

INVENTOR(S): **Rault, Sylvain**; Enguehard, Cucile;

Lancelot, Jean-Charles; Robba, Max; Atassi,

Ghanem; Pierre, Alain; **Caignard, Daniel-Henri**

; **Renard, Pierre**

PATENT ASSIGNEE(S): Adir Et Compagnie, Fr.

SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000044572	A2	20000215	JP 1999-210412	19990726
FR 2781482	A1	20000128	FR 1998-9552	19980727
FR 2781482	B1	20010831		
NO 9903608	A	20000128	NO 1999-3608	19990723
NO 313521	B1	20021014		
EP 982308	A1	20000301	EP 1999-401892	19990726
EP 982308	B1	20030226		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 9902841	A	20000509	BR 1999-2841	19990726
US 6071945	A	20000606	US 1999-361064	19990726
NZ 336913	A	20000623	NZ 1999-336913	19990726
AT 233266	E	20030315	AT 1999-401892	19990726
CA 2279099	C	20030617	CA 1999-2279099	19990726
CA 2279099	AA	20000127		
ES 2192827	T3	20031016	ES 1999-401892	19990726
ZA 9904817	A	20000202	ZA 1999-4817	19990727
CN 1244529	A	20000216	CN 1999-110693	19990727
CN 1122669	B	20031001		
AU 9941171	A1	20000217	AU 1999-41171	19990727
AU 752505	B2	20020919		
HK 1024244	A1	20040514	HK 2000-103590	20000615

PRIORITY APPLN. INFO.:


OTHER SOURCE(S):

CASREACT 132:151677; MARPAT 132:151677

GI



I



II

AB Title compds. I (R1, R3 = H, halo, C1-6 alkyl, NO2, OH, C1-6 alkoxy, etc.; R2 = (un)substituted aryl, heteroaryl), their isomers, or pharmaceutically acceptable salts are prepared from II (prepared by successive reaction of R2CH2CN (R2 = same as I) with HCO2Et, PhSO2Cl, and methylthio glycolate) via several steps. 4-(2-Methoxyphenyl)-3-(1H-1-pyrrolyl)-2-thiophene-N-pyrrolidinocarbonyl was cyclized in the presence of POCl3 under reflux for 3 h to give 3-(2-methoxyphenyl)-8H-thieno[2,3-b]pyrrolizin-8-one showing in vivo antitumor activity against leukemic P 388.

L81 ANSWER 17 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1999:614430 HCAPLUS

DOCUMENT NUMBER: 131:307037

TITLE: Effects of S-21007, a potent 5-HT3 partial agonist, in mouse anxiety

AUTHOR(S): Delagrance, Philippe; Misslin, Rene; Seale, Thomas W.;
Pfeiffer, Bruno; Rault, Sylvain;
Renard, Pierre
CORPORATE SOURCE: Institut de Recherches Internationales Servier,
Courbevoie, 92415, Fr.
SOURCE: Zhongguo Yaoli Xuebao (1999), 20(9), 805-812
CODEN: CYLPDN; ISSN: 0253-9756
PUBLISHER: Kexue Chubanshe
DOCUMENT TYPE: Journal
LANGUAGE: English
AB AIM: To study the effect of S-21007, a 5-HT3 partial agonist in different
animal models of anxiety in mice. METHODS: S-21007 effects were evaluated
in behavior tests after i.p. and oral acute treatment or in the light/dark
test after both acute and chronic treatments. RESULTS: S-21007 presented
anxiolytic-like properties after acute administration in the light/dark
box test, the mirrored chamber test, and the elevated plus-maze at low
doses 10 ng·kg⁻¹-100 µg·kg⁻¹, 1-100 µg·kg⁻¹ and
10-100 µg·kg⁻¹, resp. In the light/dark box test, S-21007 was
active orally after acute treatment at 100 ng·kg⁻¹-10
mg·kg⁻¹ and after chronic treatment (14 d) at 1-10
µg·kg⁻¹. S-21007 was devoid of sedative or stimulatory effects.
CONCLUSION: S-21007 exhibited anxiolytic-like properties. The mechanism
of action may be a desensitization of 5-HT3 receptor or an antagonist
activity on the 5-HT3 receptors.
REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

W L81 ANSWER 18 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1998:366507 HCAPLUS
DOCUMENT NUMBER: 129:122642
TITLE: Synthesis of new pyrrolo[1,2-a]quinoxalines: potential
non-peptide glucagon receptor antagonists
AUTHOR(S): Guillon, Jean; Dallemagne, Patrick; **Pfeiffer,**
Bruno; Renard, Pierre; Manechez,
Dominique; Kervran, Alain; **Rault, Sylvain**
CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de
Normandie, Lab. de Pharmacochimie, UFR des Sciences
Pharmaceutiques, Caen, 14032, Fr.
SOURCE: European Journal of Medicinal Chemistry (1998), 33(4),
293-308
CODEN: EJMCA5; ISSN: 0223-5234
PUBLISHER: Editions Scientifiques et Medicales Elsevier
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Synthesis of new pyrrolo[1,2-a]quinoxaline derivs. was achieved starting
from various nitroanilines or o-phenylenediamines. They showed little
affinity towards glucagon receptors.
REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

W L81 ANSWER 19 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1997:320915 HCAPLUS
DOCUMENT NUMBER: 127:271
TITLE: Novel and Selective Partial Agonists of 5-HT3
Receptors. 2. Synthesis and Biological Evaluation of
Piperazinopyridopyrrolopyrazines,
Piperazinopyrroloquinoxalines, and
Piperazinopyridopyrroloquinoxalines
AUTHOR(S): Prunier, Herve; **Rault, Sylvain;**
Lancelot, Jean-Charles; Robba, Max;

**Renard, Pierre; Delagrange, Philippe;
Pfeiffer, Bruno; Caignard,
Daniel-Henri; Misslin, Rene; Guardiola-Lemaitre,
Beatrice; Hamon, Michel**

CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de
Normandie, Universite de Caen, Caen, 14032, Fr.

SOURCE: Journal of Medicinal Chemistry (1997), 40(12),
1808-1819

CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In continuation of our previous work on piperazinopyrrolothienopyrazine
derivs., three series of piperazinopyridopyrrolopyrazines,
piperazinopyrroloquinoxalines, and piperazinopyridopyrroloquinoxalines
were prepared and evaluated as 5-HT₃ receptor ligands. The chemical
modifications performed within these new series led to structure-activity
relationships regarding both high affinity and selectivity for the 5-HT₃
receptors that are in agreement with those established previously for the
pyrrolothienopyrazine series. The best compound (8a) obtained in these new
series is in the picomolar range of affinity for 5-HT₃ receptors with a
selectivity higher than 106. Four of the high-affinity 5-HT₃ ligands (8a,
15a,b, and 16d) were selected in both the pyridopyrrolopyrazine and the
pyrroloquinoxaline series and were characterized in vitro and in vivo as
agonists or partial agonists. Compound 8a was also evaluated in the
light/dark test where it showed potential anxiolytic-like activity at very
low doses per os.

REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

10 L81 ANSWER 20 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1996:241880 HCAPLUS

DOCUMENT NUMBER: 124:289475

TITLE: Novel Selective and Partial Agonists of 5-HT₃
Receptors. Part 1. Synthesis and Biological Evaluation
of Piperazinopyrrolothienopyrazines

AUTHOR(S): **Rault, Sylvain; Lancelot,
Jean-Charles; Prunier, Herve; Robba, Max;
Renard, Pierre; Delagrange, Philippe;
Pfeiffer, Bruno; Caignard,
Daniel-Henri; Guardiola-Lemaitre, Beatrice;
Hamon, Michel**

CORPORATE SOURCE: Centre d'Etudes et de Recherche sur le Medicament de
Normandie, Universite de Caen, Caen, 14032, Fr.

SOURCE: Journal of Medicinal Chemistry (1996), 39(10), 2068-80
CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A series of (piperazinyl)pyrrolo[1,2-a]thieno[3,2-e]pyrazines and
(piperazinyl)pyrrolo[1,2-a]thieno[2,3-e]pyrazines was prepared and evaluated
in order to determine the necessary requirements for high affinity on the 5-HT₃
receptors and high selectivity vs. other 5-HT receptor subtypes. Various
substitutions on the piperazine and the thiophene ring of the
pyrrolothienopyrazine moieties were systematically explored as well as
replacement of the piperazine by other cyclic amines. An example compound
is 5-[4-[(4-fluorophenyl)methyl]-1-piperazinyl]pyrrolo[1,2-a]thieno[3,2-
e]pyrazine trihydrochloride. These high-affinity compds. have in common a
benzyl- or allylpiperazine substituent with no substitutions on the
thiophene ring. Five of these compds. were evaluated on the Von

Bezold-Jarisch reflex and were characterized as partial agonists. One of them, 5-[4-(phenylmethyl)-1-piperazinyl]pyrrolo[1,2-a]thieno[3,2-e]pyrazine (fumarate) was shown in vivo at very low dose a potent anxiolytic-like activity in the light/dark test.

160 L81 ANSWER 21 OF 23 HCAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1996:726818 HCAPLUS
DOCUMENT NUMBER: 126:69969
TITLE: Interaction of S 21007 with 5-HT3 receptors. In vitro and in vivo characterization
AUTHOR(S): Delagrangé, Philippe; Emerit, M. Boris; Merahi, Nacera; Abraham, Christine; Morain, Philippe; **Rault, Sylvain; Renard, Pierre; Pfeiffer, Bruno**; Guardiola-Lemaitre, Beatrice; et al.
CORPORATE SOURCE: IRIS, 6 Place des Pleiades, Courbevoie, 92415, Fr.
SOURCE: European Journal of Pharmacology (1996), 316(2/3), 195-203
CODEN: EJPHAZ; ISSN: 0014-2999
PUBLISHER: Elsevier
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The interaction of S 21007 [5-(4-benzyl piperazin-1-yl)4H pyrrolo[1,2-a]thieno[3,2-e]pyrazine] with serotonin 5-HT3 receptors was investigated using biochem., electrophysiol. and functional assays. Binding studies using membranes from N1E-115 neuroblastoma cells showed that S 21007 is a selective high affinity (IC50 = 2.8 nM) 5-HT3 receptor ligand. As expected of an agonist, S 21007 stimulated the uptake of [14C]guanidinium (EC50.apprx.10 nM) in NG 108-15 cells exposed to substance P, and this effect could be prevented by the potent 5-HT3 receptor antagonist ondansetron. In addition, like 5-HT and other 5-HT3 receptor agonists (phenylbiguanide and 3-chloro-phenylbiguanide), S 21007 (EC50 = 27 µM) produced a rapid inward current in N1E-115 cells. The 5-HT3 receptor agonist action of S 21007 was also demonstrated in urethane-anesthetized rats as this drug (120 µg/kg i.v.) triggered the Bezold-Jarisch reflex (rapid fall in heart rate), and this action could be prevented by pretreatment with the potent 5-HT3 receptor antagonist zacopride. Finally, in line with its 5-HT3 receptor agonist properties, S 21007 also triggered emesis in the ferret. Evidence for 5-HT3 receptor antagonist-like properties of S 21007 was also obtained in some of these expts. since previous exposure to this compound prevented both the 5-HT-induced current in N1E-115 cells and the Bezold-Jarisch reflex elicited by an i.v. bolus of 5-HT (30 µg/kg) in urethane-anesthetized rats. These data suggest that S 21007 is a selective 5-HT3 receptor agonist which can exhibit antagonist-like properties either by triggering a long lasting receptor desensitization or by a partial agonist activity at 5-HT3 receptors in some tissues.

160 L81 ANSWER 22 OF 23 WPIX COPYRIGHT 2006 THE THOMSON CORP on STN
ACCESSION NUMBER: 2000-149624 [14] WPIX
DOC. NO. CPI: C2000-047071
TITLE: New 8H-thieno(2,3-b)pyrrolizin-8-one derivatives useful as anticancer agents, especially for treating solid tumors.
DERWENT CLASS: B02
INVENTOR(S): ATASSI, G; **CAIGNARD, D H**; ENGUEHARD, C; **LANCELOT, J**; PIERRE, A; **RAULT, S**; **RENARD, P**; ROBBA, M; **CAIGNARD, D**; ROBBA, M; EGUHAD, C; LANSLOT, J C; LART, S; **LANCELOT, J C**

PATENT ASSIGNEE(S): (SERV-N) LES LAB SERVIER; (ADIR) ADIR & CIE
 COUNTRY COUNT: 35
 PATENT INFORMATION:

PATENT NO	KIND	DATE	WEEK	LA	PG
FR 2781482	A1	20000128	(200014)*		22
EP 982308	A1	20000301	(200016)	FR	
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT					
RO SE SI					
NO 9903608	A	20000128	(200016)		
AU 9941171	A	20000217	(200019)		
JP 2000044572	A	20000215	(200019)		12
CN 1244529	A	20000216	(200027)		
ZA 9904817	A	20000426	(200027)		27
CA 2279099	A1	20000127	(200028)	FR	
HU 9902546	A2	20000428	(200030)		
BR 9902841	A	20000509	(200033)		
US 6071945	A	20000606	(200033)		
NZ 336913	A	20000623	(200038)		
AU 752505	B	20020919	(200272)		
NO 313521	B1	20021014	(200275)		
EP 982308	B1	20030226	(200316)	FR	
R: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE					
DE 69905524	E	20030403	(200330)		
CA 2279099	C	20030617	(200347)	FR	
ES 2192827	T3	20031016	(200377)		
CN 1122669	C	20031001	(200553)		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
FR 2781482	A1	FR 1998-9552	19980727
EP 982308	A1	EP 1999-401892	19990726
NO 9903608	A	NO 1999-3608	19990723
AU 9941171	A	AU 1999-41171	19990727
JP 2000044572	A	JP 1999-210412	19990726
CN 1244529	A	CN 1999-110693	19990727
ZA 9904817	A	ZA 1999-4817	19990727
CA 2279099	A1	CA 1999-2279099	19990726
HU 9902546	A2	HU 1999-2546	19990727
BR 9902841	A	BR 1999-2841	19990726
US 6071945	A	US 1999-361064	19990726
NZ 336913	A	NZ 1999-336913	19990726
AU 752505	B	AU 1999-41171	19990727
NO 313521	B1	NO 1999-3608	19990723
EP 982308	B1	EP 1999-401892	19990726
DE 69905524	E	DE 1999-605524	19990726
		EP 1999-401892	19990726
CA 2279099	C	CA 1999-2279099	19990726
ES 2192827	T3	EP 1999-401892	19990726
CN 1122669	C	CN 1999-110693	19990727

FILING DETAILS:

PATENT NO	KIND	PATENT NO
AU 752505	B Previous Publ.	AU 9941171
NO 313521	B1 Previous Publ.	NO 9903608

DE 69905524 E Based on EP 982308
 ES 2192827 T3 Based on EP 982308

PRIORITY APPLN. INFO: FR 1998-9552 19980727

AN 2000-149624 [14] WPIX

AB FR 2781482 A UPAB: 20000320

NOVELTY - 8H-Thieno(2,3-b)pyrrolizin-8-one derivatives (I) are new.

DETAILED DESCRIPTION - 8H-Thieno(2,3-b)pyrrolizin-8-one derivatives (I) and salts are new.

R1, R3 = H, halo, 1-6C alkyl, NO2, OH, 1-6C alkoxy, 1-6C trihaloalkyl, 1-6C trihaloalkoxy, NH2, 1-6C alkylamino or di(1-6C alkyl)amino;

R2 = aryl or heteroaryl (both optionally substituted); and provided that R2 is phenyl optionally p-substituted by Br, Cl, F, OMe or OH when R1 and R3 are both H.

An INDEPENDENT CLAIM is also included for the preparation of (I).

ACTIVITY - Cytostatic. 3-(3,4-Dihydroxyphenyl)-8H-thieno(2,3-b)pyrrolizin-8-one had IC50 values of 356 and 222 nM respectively against P388 and L1210 murine leukemia cells and 122, 33 and 22 nM respectively against A549, KB-3-1 and KB-A1 human solid tumor cells.

MECHANISM OF ACTION - None given.

USE - (I) are anticancer agents especially useful for treating solid tumors.
 Dwg.0/0

L81 ANSWER 23 OF 23 WPIX COPYRIGHT 2006 THE THOMSON CORP on STN

ACCESSION NUMBER: 1994-343271 [43] WPIX

DOC. NO. CPI: C1994-156328

TITLE: Pyrrolo pyrazine derivatives having activity on 5-HT3 receptors - useful for treating anxiety, depression, migraine, pain, etc..

DERWENT CLASS: B02

INVENTOR(S): ADAM, G; DELAGRANGE, P; LANCELOT, J; PRUNIER, H; RAULT, S; RENARD, P; ROBBA, M

PATENT ASSIGNEE(S): (ADIR) ADIR & CIE

COUNTRY COUNT: 22

PATENT INFORMATION:

PATENT NO	KIND	DATE	WEEK	LA	PG
EP 623620	A1	19941109	(199443)*	FR	32
R: AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE					
FR 2704547	A1	19941104	(199444)		
AU 9461873	A	19941103	(199501)		
CA 2122290	A	19941031	(199505)	FR	
JP 06340666	A	19941213	(199509)		41
NZ 260425	A	19950427	(199522)		
ZA 9402964	A	19950426	(199522)		41
AU 671199	B	19960815	(199641)		
US 5599812	A	19970204	(199711)		10
EP 623620	B1	19980909	(199840)	FR	
R: AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE					
DE 69413112	E	19981015	(199847)		
ES 2123728	T3	19990116	(199909)		

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
EP 623620	A1	EP 1994-400881	19940425

FR 2704547	A1	FR 1993-5109	19930430
AU 9461873	A	AU 1994-61873	19940428
CA 2122290	A	CA 1994-2122290	19940427
JP 06340666	A	JP 1994-127963	19940502
NZ 260425	A	NZ 1994-260425	19940429
ZA 9402964	A	ZA 1994-2964	19940429
AU 671199	B	AU 1994-61873	19940428
US 5599812	A	US 1994-235426	19940429
EP 623620	B1	EP 1994-400881	19940425
DE 69413112	E	DE 1994-613112	19940425
		EP 1994-400881	19940425
ES 2123728	T3	EP 1994-400881	19940425

FILING DETAILS:

PATENT NO	KIND	PATENT NO
AU 671199	B Previous Publ.	AU 9461873
DE 69413112	E Based on	EP 623620
ES 2123728	T3 Based on	EP 623620

PRIORITY APPLN. INFO: FR 1993-5109 19930430

AN 1994-343271 [43] WPIX

AB EP 623620 A UPAB: 19981021

Pyrrolo pyrazine derivs. of formula (I), their optical isomers and acid addition salts are new. R1=R2-N-R3. R2 and R3 form with the N atom a group selected from (substd.) piperazine, (substd.) piperidine, (substd.) pyrrolidine, (alkyl substd.) morpholine, etc; A forms with the 2 indicated C atoms a benzo, pyrido, pyrazino, or pyrimidino ring and is opt. substd. by alkyl, OH, alkoxy, etc.

USE - The compounds act on 5-HT₃ receptors and are useful in the treatment of anxiety, depression, stress, psychoses, schizophrenia, CNS disorders, migraine, memory failure, behavioural disorders, eating disorders, alcoholism, pain, and as anti-emetics.
Dwg.0/0

ABEQ US 5599812 A UPAB: 19970313

Cpds. of formula (I) are new. R1 = N(R2)(R3); N(R2)(R3) =

piperazine,
piperidine, or
pyrrolidine,

(all opt. substd.),

morpholine (opt. substd. by one more alkyl groups),

tetrahydropyridine,

thiomorpholine,

a 5- to 12-membered azaspiro (opt. substd. by one or more alkyl or oxo groups),

a 7- to 12-membered mono- or bicyclic azacycloalkyl optionally including, in its skeleton, to 1 or 2 additional heteroatoms chosen from oxygen, sulphur, and nitrogen,

a 7- to 12-membered mono- or bicyclic azacycloalkyl, substd. by one or more alkyl or oxo groups, opt. including, in its skeleton, to 1 or 2 additional heteroatoms chosen from O, S or N,

-NH-(CH₂)_k-NH₂; k = 2-4,

and a substituted group -NH-(CH₂)_k-NH₂ in which k is as defined above,

and A forms, with the 2 carbon atoms to which it is bonded, a ring chosen from benzo, pyrido, pyrazino and pyrimidino; A being unsubstituted or substituted by one or more radicals chosen from:

alkyl,

hydroxyl,

alkoxy,
acyl,
alkoxycarbonyl,
halogen,

trifluoromethyl,

-(CH₂)_m-phenyl and -O-(CH₂)_m-phenyl in which the phenyl ring is itself unsubstituted or substituted by one or more radicals chosen from halogen, alkyl, alkoxy, hydroxyl and trifluoromethyl; and m represents 0 or 1 to 4,

-(CH₂)_m-piperazine in which the piperazine group is itself substituted or unsubstituted and m is as defined above,

provided that if A forms, with the 2 carbon atoms to which it is bonded, a benzo ring, then R₂ and R₃ cannot form, with the nitrogen atom which carries them, piperazine which is unsubstituted or substituted by alkyl, phenyl or alkyl-substituted phenyl, morpholine or aminoalkylamines,

it being understood that the term "substituted", as it relates to the piperazine, piperidine, pyrrolidine and -NH-(CH₂)_k-NH₂ groups, means that these groups can be substituted by one or more halogen, hydroxyl, oxo, R₄ radicals or -C(O)-(R₄)

with R₄ being chosen from:

alkyl,
alkoxy,

alkenyl which is unsubstituted or substituted by a phenyl which is itself unsubstituted or substituted by one or more radicals chosen from halogen, alkyl, alkoxy, hydroxyl and trifluoromethyl, -(CH₂)_n-R₅ or -(CH₂)_{n'}-C(O)-R₅,

where n = 0-5, n' = 1-5, R₅ = phenyl, benzhydryl, thienyl, pyrrolyl, pyrrolidinyl, furyl, pyrimidinyl, pyridyl, methylenedioxyphenyl, ethylenedioxyphenyl, naphthyl, quinolyl, isoquinolyl, cycloalkyl and dicycloalkylmethyl; the term "cycloalkyl" meaning a 3- to 12-membered mono- or bicyclic group, it being possible for these R₅ radicals themselves to be substituted by one or more of radicals chosen from halogen, trifluoromethyl, carboxyl, hydroxyl, alkyl and alkoxy,

and (CH₂)_{n'}-R₆ where n' is as defined above and R₆ represents a group chosen from carboxyl, alkoxycarbonyl, amino, alkylamino, dialkylamino, -SO₂N(R₇)(R₈) and -CON(R₇)(R₈) in which R₇ and R₈ represent, each independently of the other, a hydrogen atom or an alkyl group,

its optical isomers,

and its addition salts with a pharmaceutically acceptable acid or base,

it being understood that, except when otherwise specified, the terms "alkyl", "alkoxy" and "acyl" represent linear or branched groups having from 1 to 6 carbon atoms, inclusive,

and the term "alkenyl" represents a linear or branched unsaturated group having from 2 to 6 carbon atoms, inclusive.

Dwg.0/0

=> □

=> file registry

FILE 'REGISTRY' ENTERED AT 11:27:26 ON 19 JUL 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

STRUCTURE
SEARCH

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3
DICTIONARY FILE UPDATES: 18 JUL 2006 HIGHEST RN 894196-03-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when
conducting SmartSELECT searches.


REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> d stat que L42
L14 STR

X 1

Ak.....O 2



G1 [@1], [@2], [@3], [@4], [@5]

Structure attributes must be viewed using STN Express query preparation.

L16 29 SEA FILE=REGISTRY SSS FUL L14
L42 4 SEA FILE=HCAPLUS ABB=ON PLU=ON L16

=> =>

=> file hcaplus

FILE 'HCAPLUS' ENTERED AT 11:31:17 ON 19 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2006 VOL 145 ISS 4

FILE LAST UPDATED: 18 Jul 2006 (20060718/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

'OBI' IS DEFAULT SEARCH FIELD FOR 'HCAPLUS' FILE

=> d que nos L42

L14 STR

L16 29 SEA FILE=REGISTRY SSS FUL L14

L42 4 SEA FILE=HCAPLUS ABB=ON PLU=ON L16

=> s L42 not L79

L83 3 L42 NOT L79

printed with author search

=> file babs

FILE 'BABS' ENTERED AT 11:32:30 ON 19 JUL 2006

COPYRIGHT (c) 2006 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE LAST UPDATED: 15 JUN 2006 <20060615/UP>

FILE COVERS 1980 TO DATE.

=> d stat que L41

L41 1 SEA FILE=BABS ABB=ON PLU=ON 5564377/BABSAN

=> file beilstein

FILE 'BEILSTEIN' ENTERED AT 11:32:55 ON 19 JUL 2006

COPYRIGHT (c) 2006 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE LAST UPDATED ON JUNE 16, 2006

FILE COVERS 1771 TO 2006.

*** FILE CONTAINS 9,606,495 SUBSTANCES ***

>>>PLEASE NOTE: Reaction Data and substance data are stored in

separate documents and can not be searched together in one query. Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a compounds with available reaction information by combining with PRE/FA, REA/FA or more generally with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be searched as reaction partner BRNs Reactant BRN (RX.RBRN) or Product BRN (RX.PBRN).<<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

 * PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
 * SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
 * ARE BASED ON THE HIGHEST PRICE CATEGORY. THEREFORE; THESE *
 * ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
 * FOR PRICE INFORMATION SEE HELP COST *


NEW

* PATENT NUMBERS (PN) AND BABS ACCESSION NUMBERS (BABSAN) CAN NOW BE SEARCHED, SELECTED AND TRANSFERRED.
 * NEW DISPLAY FORMATS ALLREF, ALLP AND BABSAN SHOW ALL REFERENCES, ALL PATENT REFERENCES, OR ALL BABS ACCESSION NUMBERS FOR A COMPOUND AT A GLANCE.

=> d stat que L22
 L5 STR

X 1

Ak O 2




G1 [@1], [@2], [@3], [@4], [@5]

Structure attributes must be viewed using STN Express query preparation.
 L7 16 SEA FILE=REGISTRY SSS FUL L5
 L14 STR

X 1

Ak.....O 2



G1 [@1], [@2], [@3], [@4], [@5]

Structure attributes must be viewed using STN Express query preparation.

L16 29 SEA FILE=REGISTRY SSS FUL L14
 L17 13 SEA FILE=REGISTRY ABB=ON PLU=ON L16 NOT L7
 L18 3 SEA FILE=CAPLUS ABB=ON PLU=ON L17
 L21 8 SEA FILE=BEILSTEIN SSS FUL L14
 L22 1 SEA FILE=BEILSTEIN ABB=ON PLU=ON L21 NOT L18

→ compounds that
 have CAPLUS
 references

=> file wpix

FILE 'WPIX' ENTERED AT 11:33:26 ON 19 JUL 2006

COPYRIGHT (C) 2006 THE THOMSON CORPORATION

FILE LAST UPDATED: 14 JUL 2006 <20060714/UP>
 MOST RECENT DERWENT UPDATE: 200645 <200645/DW>
 DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,
 PLEASE VISIT:
http://www.stn-international.de/training_center/patents/stn_guide.pdf <

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES, SEE
<http://scientific.thomson.com/support/patents/coverage/latestupdates/>

>>> PLEASE BE AWARE OF THE NEW IPC REFORM IN 2006, SEE
http://www.stn-international.de/stndatabases/details/ipc_reform.html and
<http://scientific.thomson.com/media/scpdf/ipcrdwpi.pdf> <<<


>>> FOR FURTHER DETAILS ON THE FORTHCOMING DERWENT WORLD PATENTS
 INDEX ENHANCEMENTS PLEASE VISIT:
http://www.stn-international.de/stndatabases/details/dwpi_r.html <<<
 'BIX' IS DEFAULT SEARCH FIELD FOR 'WPIX' FILE

=> d stat que L39

L14 STR

X 1

Ak.....O 2



G1 [@1], [@2], [@3], [@4], [@5]

Structure attributes must be viewed using STN Express query preparation.

L35 16 SEA FILE=WPIX SSS FUL L14
 L36 2 SEA FILE=WPIX ABB=ON PLU=ON L35/DCR
 L37 2 SEA FILE=WPIX ABB=ON PLU=ON (RAEA00/DCN OR RAEA01/DCN OR
 RAEA02/DCN OR RAEA03/DCN OR RAEA04/DCN OR RAEA05/DCN OR
 RAEA07/DCN OR RAEA08/DCN OR RAE9ZS/DCN OR RAE9ZT/DCN OR
 RAE9ZU/DCN OR RAE9ZV/DCN OR RAE9ZW/DCN OR RAE9ZX/DCN OR
 RAE9ZY/DCN OR RAFSCY/DCN)
 L38 2 SEA FILE=WPIX ABB=ON PLU=ON (902542-0-0-0/DCRE OR 902543-0-0-0/DCRE OR 902544-1-0-0/DCRE OR 902544-2-0-0/DCRE OR 902546-0-0-0/DCRE OR 902547-0-0-0/DCRE OR 902548-1-0-0/DCRE OR 902548-2-0-0/DCRE OR 902550-1-0-0/DCRE OR 902551-0-0-0/DCRE OR 902552-1-0-0/DCRE OR 902552-2-0-0/DCRE OR 902553-0-0-0/DCRE OR 902557-0-0-0/DCRE OR 902558-0-0-0/DCRE OR 976436-1-0-0/DCRE)
 L39 2 SEA FILE=WPIX ABB=ON PLU=ON (L36 OR L37 OR L38)

=> s L39 not L80

L84

1 L39 NOT

L80 → printed with author search

=> file marpat

FILE 'MARPAT' ENTERED AT 11:33:55 ON 19 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

FILE CONTENT: 1961-PRESENT VOL 145 ISS 1 (20060714/ED)

SOME MARPAT RECORDS ARE DERIVED FROM INPI DATA FOR 1961-1987

MOST RECENT CITATIONS FOR PATENTS FROM MAJOR ISSUING AGENCIES
 (COVERAGE TO THESE DATES IS NOT COMPLETE):

US 2006118302 08 JUN 2006

DE 102004053653 04 MAY 2006
EP 1653548 03 MAY 2006
JP 2006112980 27 APR 2006
WO 2006053912 26 MAY 2006
GB 2419594 03 MAY 2006
FR 2877004 28 APR 2006
RU 2275374 27 APR 2006
CA 2518664 10 MAR 2006

Expanded G-group definition display now available.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

=> d stat que L30
L5 STR


X 1

Ak O 2

Cb 3

N 4

Hy 5




G1 [@1], [@2], [@3], [@4], [@5]

note:
← search query for
MARPAT is narrower
than for other
files

Structure attributes must be viewed using STN Express query preparation.
L14 STR

X 1

Ak O 2



G1 [@1], [@2], [@3], [@4], [@5]

Structure attributes must be viewed using STN Express query preparation.

L28 49 SEA FILE=MARPAT SSS FUL L14

L30 24 SEA FILE=MARPAT SUB=L28 SSS FUL L5

100.0% PROCESSED 30 ITERATIONS

24 ANSWERS

SEARCH TIME: 00.00.01

=> s L30 not L68

L85 23 L30 NOT L68

printed with author search

=> => dup rem L83 L84 L41 L85 L22

DUPLICATE IS NOT AVAILABLE IN 'BEILSTEIN'.

ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE

FILE 'HCAPLUS' ENTERED AT 11:35:09 ON 19 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 11:35:09 ON 19 JUL 2006

COPYRIGHT (C) 2006 THE THOMSON CORPORATION

FILE 'BABS' ENTERED AT 11:35:09 ON 19 JUL 2006

COPYRIGHT (c) 2006 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften
licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE 'MARPAT' ENTERED AT 11:35:09 ON 19 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

FILE 'BEILSTEIN' ENTERED AT 11:35:09 ON 19 JUL 2006

COPYRIGHT (c) 2006 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften

licensed to Beilstein GmbH and MDL Information Systems GmbH

PROCESSING COMPLETED FOR L83

PROCESSING COMPLETED FOR L84

PROCESSING COMPLETED FOR L41

PROCESSING COMPLETED FOR L85

PROCESSING COMPLETED FOR L22

L86 26 DUP REM L83 L84 L41 L85 L22 (3 DUPLICATES REMOVED)

ANSWERS '1-3' FROM FILE HCAPLUS

ANSWERS '4-25' FROM FILE MARPAT

ANSWER '26' FROM FILE BEILSTEIN

=> d ibib abs hitstr L86 1-3; d ibib abs hit L86 4-25; d ide allref L86 26

L86 ANSWER 1 OF 26 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 2004:857326 HCAPLUS

DOCUMENT NUMBER: 141:309639

TITLE: Dipeptidyl peptidase inhibitors

INVENTOR(S): Feng, Jun; Gwaltney, Stephen L.; Kaldor, Stephen W.;
Stafford, Jeffrey A.; Wallace, Michael B.; Zhang,
Zhiyuan

PATENT ASSIGNEE(S): Syrrx, Inc., USA

SOURCE: PCT Int. Appl., 244 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004087053	A2	20041014	WO 2004-US9217	20040324
WO 2004087053	C2	20041111		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2518465	AA	20041014	CA 2004-2518465	20040324
US 2004242568	A1	20041202	US 2004-809636	20040324
US 2004242566	A1	20041202	US 2004-809638	20040324
US 2004259870	A1	20041223	US 2004-809637	20040324
US 2005004117	A1	20050106	US 2004-809635	20040324
EP 1608317	A2	20051228	EP 2004-758366	20040324
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK			
PRIORITY APPLN. INFO.:			US 2003-457785P	P 20030325
			WO 2004-US9217	W 20040324
OTHER SOURCE(S):	MARPAT 141:309639			
GI				



AB Dipeptidyl peptidase IV inhibitors I [Q = CO, SO, SO₂, C:NR₅; R₁ = ZR₆; Z = moiety providing 1-6 atom separation between R₆ and ring; R₂ = (substituted)3-7-membered ring; R₃,R₄ = taken together form a (substituted)5-6-membered ring; R₅ = H, (substituted)alkyl, cycloalkyl, etc.; R₆ = (substituted)C3-7-cycloalkyl or aryl] are disclosed. Thus, 2-[2-(3-aminopiperidin-1-yl)-6,7-dimethoxy-4-oxo-4H-quinazolin-3-ylmethyl]benzonitrile (I; R₁ = 2-cyanophenylmethyl; R₂ = 3-aminopiperidin-1-yl; R₃,R₄ = dimethoxyphenyl) was synthesized. This compound exhibited enhanced stability in rat liver microsomes.

IT 769157-67-7P 769157-79-1P 769157-80-4P

RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(dipeptidyl peptidase inhibitors)

RN 769157-67-7 HCAPLUS


CN Benzonitrile, 2-[[2-[(3R)-3-amino-1-piperidinyl]-6-methoxy-4-oxopyrido[3,4-d]pyrimidin-3(4H)-yl]methyl]-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

CRN 769157-66-6

CMF C21 H22 N6 O2


Absolute stereochemistry.



CM 2

CRN 76-05-1

CMF C2 H F3 O2




RN 769157-79-1 HCAPLUS
 CN Benzonitrile, 2-[[2-[(3R)-3-amino-1-piperidinyl]-6-chloro-4-oxopyrido[3,4-d]pyrimidin-3(4H)-yl]methyl]-, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1


CRN 769157-78-0
 CMF C20 H19 Cl N6 O

Absolute stereochemistry.




CM 2

CRN 76-05-1
 CMF C2 H F3 O2




RN 769157-80-4 HCAPLUS
 CN Benzonitrile, 2-[[2-[(3R)-3-amino-1-piperidinyl]-4-oxo-6-(1-pyrrolidinyl)pyrido[3,4-d]pyrimidin-3(4H)-yl]methyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




L86 ANSWER 2 OF 26 HCAPLUS COPYRIGHT 2006 ACS on STN DUPLICATE 2
 ACCESSION NUMBER: 1985:113414 HCAPLUS
 DOCUMENT NUMBER: 102:113414
 TITLE: Synthesis and reactions of 2,3-diamino-4(3H)-pyrimidinones and 3-amino-2-hydrazino-4(3H)-pyrimidinones. I
 AUTHOR(S): Hlavka, Joseph J.; Bitha, Panayota; Lin, Yang I.; Strohmeyer, Timothy
 CORPORATE SOURCE: Med. Res. Div., Am. Cyanamid Co., Pearl River, NY, 10965, USA
 SOURCE: Journal of Heterocyclic Chemistry (1984), 21(5), 1537-41
 CODEN: JHTCAD; ISSN: 0022-152X
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 102:113414
 GI




AB A series of new 2,3-diamino-4-pyrimidinones and 3-amino-2-hydrazino-4-pyrimidinones were synthesized by the reactions of β -ketoesters with amino or diaminoguanidines. E.g., refluxing aminoguanidine bicarbonate with $\text{CF}_3\text{COCH}_2\text{CO}_2\text{Et}$ in BuOH gave 45% pyrimidine I.
 IT 95095-61-7P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)
 RN 95095-61-7 HCAPLUS
 CN Pyrido[3,4-d]pyrimidin-4(3H)-one, 2,3-diamino-5,6,7,8-tetrahydro-7-(phenylmethyl)- (9CI) (CA INDEX NAME)



L86 ANSWER 3 OF 26 HCAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1979:593248 HCAPLUS
 DOCUMENT NUMBER: 91:193248
 TITLE: Polyaza steroids. II. 1,2,4-triazolo[3',4':2,3]pyrimido[4,5-c]quinolin-11(12H)ones, imidazo[2',1':2,3]pyrimido[4,5-c]quinolin-11(12H)ones and 2,3-dihydroimidazo[2',1':2,3]pyrimido[4,5-c]quinolin-11(12H)ones
 AUTHOR(S): Lalezari, I.; Sadeghi-Milani, S.
 CORPORATE SOURCE: Coll. Pharm., Tehran Univ., Teheran, Iran
 SOURCE: Journal of Heterocyclic Chemistry (1979), 16(4), 707-10
 CODEN: JHTCAD; ISSN: 0022-152X
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



I




II

AB 2-Substituted quinoline-3,4-dicarboxylic acids were converted in several steps to pyrimido[4,5-c]quinolinones I (R = Me, Ph; R1 = e.g. MeS, NHNH2, NHNHCOH), which were further converted to title polyazasteroids II (X = N, R = Me, Pr, Ph; R2 = H, Me, Et) and II (X = CH; same R, R2). Treating I (R1 = NHNH2) with HONO gave only I (R1 = N3) and not the tetrazolo analog of II (X = N).

IT **71881-56-6P**
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and condensation of, with chloroacetaldehyde)

RN 71881-56-6 HCAPLUS

CN Pyrimido[4,5-c]quinolin-1(2H)-one, 3-amino-2-methyl-5-phenyl- (9CI) (CA INDEX NAME)




IT 71881-50-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and cyclization. of)

RN 71881-50-0 HCAPLUS

CN Hydrazinecarboxaldehyde, 2-(1,2-dihydro-2-methyl-1-oxo-5-phenylpyrimido[4,5-c]quinolin-3-yl)- (9CI) (CA INDEX NAME)




IT 71881-59-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation and cyclodehydration of)

RN 71881-59-9 HCAPLUS

CN Pyrimido[4,5-c]quinolin-1(2H)-one, 3-[(2-hydroxyethyl)amino]-2-methyl-5-phenyl- (9CI) (CA INDEX NAME)




IT 71881-53-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation and reduction of)

RN 71881-53-3 HCAPLUS

CN Pyrimido[4,5-c]quinolin-1(2H)-one, 3-azido-2-methyl-5-phenyl- (9CI) (CA INDEX NAME)




IT 71881-78-2P 71881-79-3P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


RN 71881-78-2 HCAPLUS

CN Acetic acid, 2-(1,2-dihydro-2-methyl-1-oxo-5-phenylpyrimido[4,5-c]quinolin-3-yl)hydrazide (9CI) (CA INDEX NAME)



RN 71881-79-3 HCAPLUS

CN Propanoic acid, 2-(1,2-dihydro-2-methyl-1-oxo-5-phenylpyrimido[4,5-c]quinolin-3-yl)hydrazide (9CI) (CA INDEX NAME)




IT 71881-43-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation, acylation and reaction with nitrous acid)

RN 71881-43-1 HCAPLUS

CN Pyrimido[4,5-c]quinoline-1,3(2H,4H)-dione, 2-methyl-5-phenyl-, 3-hydrazone (9CI) (CA INDEX NAME)



46


L86 ANSWER 4 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 144:292772 MARPAT
 TITLE: Preparation of quinazolinone derivatives as B-Raf inhibitors
 INVENTOR(S): Aquila, Brian; Dakin, Les; Ezhuthachan, Jayachandran; Lee, Stephen; Lyne, Paul; Pontz, Timothy; Zheng, Xiaolan
 PATENT ASSIGNEE(S): Astrazeneca AB, Swed.; Astrazeneca UK Limited
 SOURCE: PCT Int. Appl., 112 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006024834	A1	20060309	WO 2005-GB3334	20050826
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.:

US 2004-605762P 20040831

GI




AB The title compds. I [ring A carbocyclyl, heterocyclyl; R1 = halo, nitro, cyano, etc.; n = 0-4; R2 = H, halo, nitro, etc.; X = NR15, O; one of A, E, G and J is C which is attached to X, and the other three are selected from CR16 or N; R3, R16 = H, halo, nitro, etc.; R4, R5, R15 = H, alkyl, alkanoyl, etc.] which possess B-Raf inhibitory activity and are accordingly useful for their anti cancer activity and thus in methods of treatment of the human or animal body, were prepared Thus, reacting N-(3-amino-4-methylphenyl)-3-(1-cyano-1-methylethyl)benzamide with 6-bromo-3-methylquinazolin-4(3H)-one (preps. of the reactants are given) in the presence of Pd2(dba)3, BINAP and tert-BuONa in PhMe afforded 59% 3-(1-cyano-1-methylethyl)-N-{4-methyl-3-[(3-methyl-4-oxo-3,4-dihydroquinazolin-6-yl)amino]phenyl}benzamide. The compds. I showed IC50 of <30 μ M when tested in B-Raf in vitro ELISA assay (specific IC50 values were given for representative compds. I). The invention also relates to processes for the manufacture of compds. I, to pharmaceutical compns. containing them and to their use in the manufacture of medicaments of use in the production of an anti-cancer effect in a warm blooded animal such as man.

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1



G29—G10

G1 = carbocycle <containing 3-12 C, mono- or bicyclic>
 (opt. substd. by (up to 4) G2) /
 heterocycle <containing 4-12 atoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 mono- or bicyclic> (opt. substd. by (up to 4) G2) /
 heterocycle <containing 1 or more N,
 attached through 1 or more N> (opt. substd. by 1 or more G8)
 / (Specifically claimed: 392 / thienyl / pyridyl /
 thiazolyl / isoxazolyl / furyl / 196 / 203 / 207 / 217 /
 227 / pyrazolyl (opt. substd. by 1 or more G22) / indolyl /
 241 / 251 / 257 / 261 / 271 / 281 / 289 / 299 / 311 / 321 /
 331 / 341 / pyrimidinyl / 410 / 425 / 435 / 446 / 457)




G2 = R / F / Cl / Br / I / NO₂ / CN / OH / NH₂ / CO₂H /
 CONH₂ / SH / SO₂NH₂ / alkyl <containing 1-6 C>
 (opt. substd.) / alkenyl <containing 2-6 C> (opt. substd.) /
 alkynyl <containing 2-6 C> (opt. substd.) /
 alkoxy <containing 1-6 C> (opt. substd.) /
 CHO (opt. substd.) / alkylcarbonyl <containing 1-5 C>
 (opt. substd.) / OCHO (opt. substd.) /
 alkylcarbonyloxy <containing 1-5 C> (opt. substd.) /
 alkylamino <containing 1-6 C> (opt. substd.) /


dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
 NHCHO / alkylcarbonylamino <containing 1-5 C>
 (opt. substd.) / alkylaminocarbonyl <containing 1-6 C>
 (opt. substd.) / dialkylaminocarbonyl <each alkyl containing
 1-6 C> (opt. substd.) / 183 / alkoxycarbonyl <containing 1-6
 C> (opt. substd.) / alkylaminosulfonyl <containing 1-6 C>
 (opt. substd.) / dialkylaminosulfonyl <each alkyl containing
 1-6 C> (opt. substd.) / alkylsulfonylamino <containing 1-6 C>
 (opt. substd.) / 17 / 21 / 24 / 176 / 179 /
 carbocycle <containing 3-12 C, mono- or bicyclic>
 (opt. substd.) / heterocycle <containing 4-12 atoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>
 (opt. substd.) / heterocycle <containing 1 or more N,
 attached through 1 or more N> (opt. substd. by 1 or more G8)



G3 = O / NH / 19 / C(O) / S / S(O) / SO2



G4 = alkoxycarbonyl <containing 1-6 C> (opt. substd.) /
 alkyl <containing 1-6 C> (opt. substd.) /
 G5 = carbocycle <containing 3-12 C, mono- or bicyclic>
 (opt. substd.) / heterocycle <containing 4-12 atoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>
 (opt. substd.) /
 G6 = C(O) / SO2
 G7 = NH / 27











G8 = alkyl <containing 1-6 C> / CHO /
 alkylcarbonyl <containing 1-5 C> /
 alkylsulfonyl <containing 1-6 C> /
 alkoxycarbonyl <containing 1-6 C> / CONH2 /
 alkylaminocarbonyl <containing 1-6 C> /
 dialkylaminocarbonyl <each alkyl containing 1-6 C> / CH2Ph /
 CO2CH2Ph / CPh / SO2Ph
 G9 = H / F / Cl / Br / I / NO2 / CN / OH / NH2 / CO2H /
 CONH2 / SH / SO2NH2 / alkyl <containing 1-6 C>
 (opt. substd.) / alkenyl <containing 2-6 C> (opt. substd.) /
 alkynyl <containing 2-6 C> (opt. substd.) /
 alkoxy <containing 1-6 C> (opt. substd.) /
 CHO (opt. substd.) / alkylcarbonyl <containing 1-5 C>

(opt. substd.) / OCHO (opt. substd.) /
 alkylcarbonyloxy <containing 1-5 C> (opt. substd.) /
 alkylamino <containing 1-6 C> (opt. substd.) /
 dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
 NHCHO / alkylcarbonylamino <containing 1-5 C>
 (opt. substd.) / alkylaminocarbonyl <containing 1-6 C>
 (opt. substd.) / dialkylaminocarbonyl <each alkyl containing
 1-6 C> (opt. substd.) / 185 / alkoxy carbonyl <containing 1-6
 C> (opt. substd.) / alkylaminosulfonyl <containing 1-6 C>
 (opt. substd.) / dialkylaminosulfonyl <each alkyl containing
 1-6 C> (opt. substd.) / alkylsulfonylamino <containing 1-6 C>
 (opt. substd.) / carbocycle <containing 3-12 C,
 mono- or bicyclic> (opt. substd.) /
 heterocycle <containing 4-12 atoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 mono- or bicyclic> (opt. substd.) / 37 / 39 / 42 /
 heterocycle <containing 1 or more N,
 attached through 1 or more N> (opt. substd. by 1 or more G8)




G3—G5 G6—G7—G5 G7—G6—G5 G20—G21
 37 39 42 185

G10 = 47 / 56 / 65 / 80 / 103 / 116 / 129 / 148








G11 = N / CH (opt. substd.)
 G12 = H / alkyl <containing 1-6 C> (opt. substd.) /
 CHO (opt. substd.) / alkylcarbonyl <containing 1-5 C>
 (opt. substd.) / alkylsulfonyl <containing 1-6 C>
 (opt. substd.) / alkoxy carbonyl <containing 1-6 C>
 (opt. substd.) / CONH2 / carbocycle <containing 3-12 C,
 mono- or bicyclic> (opt. substd.) /


heterocycle <containing 4-12 atoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
mono- or bicyclic> (opt. substd.) /
alkylaminocarbonyl <containing 1-6 C> (opt. substd.) /
dialkylaminocarbonyl <each alkyl containing 1-6 C>
(opt. substd.) / (Specifically claimed: Me / 346 / Et /
CH₂CH₂OH / 373 / 381 / cyclopropyl)




G13 = H / F / Cl / Br / I / NO₂ / CN / OH / NH₂ / CO₂H /
CONH₂ / SH / SO₂NH₂ / alkyl <containing 1-6 C>
(opt. substd.) / alkenyl <containing 2-6 C> (opt. substd.) /
alkynyl <containing 2-6 C> (opt. substd.) /
alkoxy <containing 1-6 C> (opt. substd.) /
CHO (opt. substd.) / alkylcarbonyl <containing 1-5 C>
(opt. substd.) / OCHO (opt. substd.) /
alkylcarbonyloxy <containing 1-5 C> (opt. substd.) /
alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
NHCHO / alkylcarbonylamino <containing 1-5 C>
(opt. substd.) / alkylaminocarbonyl <containing 1-6 C>
(opt. substd.) / dialkylaminocarbonyl <each alkyl containing
1-6 C> (opt. substd.) / 187 / alkoxycarbonyl <containing 1-6
C> (opt. substd.) / alkylaminosulfonyl <containing 1-6 C>
(opt. substd.) / dialkylaminosulfonyl <each alkyl containing
1-6 C> (opt. substd.) / alkylsulfonylamino <containing 1-6 C>
(opt. substd.) / 157 / 159 / 162 /
heterocycle <containing 1 or more N,
attached through 1 or more N> (opt. substd. by 1 or more G8)
/ (Specifically claimed: Me / 342 / NMe₂ / SMe)



G14 = H / 397 / 402 / Bu-t




G16 = NH / 174 / O






G17 = alkyl <containing 1-6 C> (opt. substd.) /
CHO (opt. substd.) / alkylcarbonyl <containing 1-5 C>
(opt. substd.) / alkylsulfonyl <containing 1-6 C>
(opt. substd.) / alkoxycarbonyl <containing 1-6 C>
(opt. substd.) / CONH₂ (opt. substd.) /


carbocycle <containing 3-12 C, mono- or bicyclic>
 (opt. substd.) / heterocycle <containing 4-12 atoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>
 (opt. substd.) / alkylaminocarbonyl <containing 1-6 C>
 (opt. substd.) / dialkylaminocarbonyl <each alkyl containing
 1-6 C> (opt. substd.) / (Specifically claimed: Me / 386)




G18 = alkoxy <containing 1-6 C> (opt. substd.)
 G19 = alkyl <containing 1-6 C> (opt. substd.)
 G20 = S / S(O) / SO₂
 G21 = alkyl <containing 1-6 C> (opt. substd.)
 G22 = alkyl <containing 1-6 C>
 G23 = 352 / cyclopropyl / 359 / 367




G24 = H / 369




G25 = NH₂ / 377 / morpholino




G26 = F / 417



G27 = F / H
 G28 = 7 / NH₂



G29 = 10 / R <"displaceable group"> / NH₂ / 466 / OH




G30 = cyclopropyl / alkyl <containing 2-6 C>
(opt. substd.)

Patent location: claim 1
 Note: or pharmaceutically acceptable salts and S-oxides
 Note: additional substitution also claimed
 Note: additional oxo substitution also claimed
 Note: substitution is restricted
 Note: also incorporates claim 13, formulae II, V, and VII

L86 ANSWER 5 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 144:274309 MARPAT
 TITLE: Preparation of heteroaryl amines as antibacterial agents
 INVENTOR(S): Pierau, Sabine; Dale, Glenn
 PATENT ASSIGNEE(S): Morphochem Aktiengesellschaft fuer Kombinatorische Chemie, Germany
 SOURCE: PCT Int. Appl., 170 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006021448	A1	20060302	WO 2005-EP9204	20050825
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
DE 102004041163	A1	20060302	DE 2004-10200404116320040825	
PRIORITY APPLN. INFO.:			DE 2004-10200404116320040825	
GI				



AB The title heteroaryl amines I [wherein X1-X6 = independently N or (un)substituted CH; A = -CH₂-CO-, -CH₂-SO₂-, -NH-SO₂-, -CO-NH-, etc.; R1 = H, OH, NH₂, halo, (hetero)alkyl, etc.; R3 = (un)substituted piperidinyl, cyclohexyl, morpholino, pyrrolidino, etc.], or pharmacol. acceptable salts, solvates, hydrates, or formulations thereof were prepared as antibacterial agents. For example, II was prepared in a multi-step synthesis. II showed an MIC ≤ 2 µg/mL against at least two organisms.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1




G1 = H / F / Cl / Br / I / **NH₂** / SH /
 alkyl <containing 1-20 C> (opt. substd. by G3) /
 R <"heteroalkyl", containing zero or more N, zero or more O, zero or more P, zero or more S, zero or more Se, zero or more Si (no other heteroatoms), 0 or more double bonds, 0 or more triple bonds, no rings> /
 14 / carbocycle <containing 3-14 C, 0 or more double bonds, mono- or polycyclic> (opt. substd. by G4) /
 heterocycle <containing 3-14 atoms, zero or more N, zero or more O, zero or more P, zero or more S, zero or more Se, zero or more Si (no other heteroatoms), 0 or more double bonds, mono- or polycyclic> /
 (Specifically claimed: OMe)












G2 = alkyl <containing 1-20 C> (opt. substd. by G3) /
 R <"heteroalkyl", containing zero or more N, zero or more O, zero or more P, zero or more S, zero or more Se, zero or more Si (no other heteroatoms), 0 or more double bonds, 0 or more triple bonds, no rings> /
 carbocycle <containing 3-14 C, 0 or more double bonds,

- mono- or polycyclic> (opt. substd. by G4) /
heterocycle <containing 3-14 atoms, zero or more N,
zero or more O, zero or more P, zero or more S,
zero or more Se, zero or more Si (no other heteroatoms),
0 or more double bonds, mono- or polycyclic>
G3 = carbocycle <containing 3-14 C,
0 or more double bonds, mono- or polycyclic>
G4 = alkyl <containing 1-20 C> /
alkenyl <containing 2-20 C> / alkynyl <containing 2-20 C> /
R <"heteroalkyl", containing zero or more B, zero or more N,
zero or more O, zero or more P, zero or more S,
zero or more Se, zero or more Si (no other heteroatoms),
0 or more double bonds, 0 or more triple bonds, no rings>
G5 = N / 16



- G6 = H / F / Cl / Br / I / OH / NH2 /
alkyl <containing 1-20 C> / alkenyl <containing 2-20 C> /
alkynyl <containing 2-20 C> / R <"heteroalkyl",
containing zero or more B, zero or more N, zero or more O,
zero or more P, zero or more S, zero or more Se,
zero or more Si (no other heteroatoms),
0 or more double bonds, 0 or more triple bonds, no rings>
G7 = 24-11 20-18 / 30-11 26-18 / 36-11 32-18 /
39-11 43-18 / 49-11 46-18 / 51-11 53-18 / 56-11 58-18 /
64-11 61-18 / 72-11 69-18





- G8 = alkyl <containing 1-20 C> (opt. substd. by G3) /
alkyl <containing 1-20 C> (substd. by 1 or more G9) /

alkenyl <containing 2-20 C> (opt. substd. by 1 or more G9) /
 alkynyl <containing 2-20 C> (opt. substd. by 1 or more G9) /
 R <"heteroalkyl", containing zero or more B, zero or more N,
 zero or more O, zero or more P, zero or more S,
 zero or more Se, zero or more Si (no other heteroatoms),
 0 or more double bonds, 0 or more triple bonds, no rings> /
 aryl <containing 6-14 C, mono- or polycyclic> /
 heteroaryl <containing 5-14 atoms, zero or more N,
 zero or more O, zero or more P,
 zero or more S (no other heteroatoms), mono- or polycyclic> /
 carbocycle <containing 3-14 C, 0 or more double bonds,
 mono- or polycyclic> (opt. substd. by G4) /
 carbocycle <containing 3-14 C, 0 or more double bonds,
 mono- or polycyclic> (substd. by 1 or more G9) /
 heterocycle <containing 3-14 atoms, zero or more N,
 zero or more O, zero or more P, zero or more S,
 zero or more Se, zero or more Si (no other heteroatoms),
 0 or more double bonds, mono- or polycyclic> /
 (Specifically claimed: 96)



G23-G24
 96 97

G9 = aryl <containing 6-14 C, mono- or polycyclic> /
 heteroaryl <containing 5-14 atoms, zero or more N,
 zero or more O, zero or more P,
 zero or more S (no other heteroatoms), mono- or polycyclic>
 G10 = CH (opt. substd.) / N
 G11 = CH / N
 G12 = 75-7 76-12 / 77-7 78-12 / 80-7 79-12 /
 83-7 84-12 / 85-7 86-12 / 87-7 88-12 / 91-7 92-12 /
 (Specifically claimed: 93-7 94-12)




G13 = NH (opt. substd.) / CH2 (opt. substd.)
 G14 = CH2 (opt. substd.)
 G15 = OH (opt. substd.)
 G16 = H / R
 G17 = C(O) / CH2 (opt. substd.)
 G18 = NH (opt. substd.)
 G19 = CH2 (opt. substd.)
 G20 = O / S
 G21 = alkyl <containing 1-4 C>

G22 = O / CH₂ (opt. substd.)
 G23 = alkylene / alkenylene / alkynylene / NH /
 R <"heteroalkylene"> / 98-12 99-97 / 101-12 103-97 /
 104-12 106-97 / 107-12 110-97 / 111-12 112-97 /
 113-12 114-97



G24 = alkyl <containing 1-20 C> (substd. by G3) /
 alkyl <containing 1-20 C> (substd. by 1 or more G9) /
 alkenyl <containing 2-20 C> (substd. by 1 or more G9) /
 alkynyl <containing 2-20 C> (substd. by 1 or more G9) /
 R <"heteroalkyl", containing zero or more B, zero or more N,
 zero or more O, zero or more P, zero or more S,
 zero or more Se, zero or more Si (no other heteroatoms),
 0 or more double bonds, 0 or more triple bonds, no rings> /
 aryl <containing 6-14 C, mono- or polycyclic> /
 heteroaryl <containing 5-14 atoms, zero or more N,
 zero or more O, zero or more P,
 zero or more S (no other heteroatoms), mono- or polycyclic> /
 carbocycle <containing 3-14 C, 0 or more double bonds,
 mono- or polycyclic> (opt. substd. by G4) /
 heterocycle <containing 3-14 atoms, zero or more N,
 zero or more O, zero or more P, zero or more S,
 zero or more Se, zero or more Si (no other heteroatoms),
 0 or more double bonds, mono- or polycyclic> / 121 / 124 /
 134 / 144 / 197 / 174 / 177



G25 = CH / N
 G26 = O / S
 G27 = H / R
 Patent location:
 Note:

claim 1
 or pharmacologically acceptable salts, solvates,
 hydrates or pharmacologically acceptable


Note: formulation
additional substitution also claimed

Pass
Index 7

*L86- ANSWER 6 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 144:150378 MARPAT
 TITLE: Preparation of pyrido[2,3-d]pyrimidine-2,4-diones and
 related compounds as selective dipeptidyl peptidase
 inhibitors
 INVENTOR(S): Feng, Jun; Gwaltney, Stephen L.; Lam, Betty; Zhang,
 Zhiyuan
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 55 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2006014764	A1	20060119	US 2005-183335	20050715
WO 2006019965	A2	20060223	WO 2005-US25070	20050714
WO 2006019965	A3	20060406		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM WO 2006020017 A2 20060223 WO 2005-US25153 20050715 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRIORITY APPLN. INFO.: US 2004-588577P 20040716
 GI




AB Pyrido[2,3-d]pyrimidine-2,4-diones and related compds. (shown as I; variables defined below; e.g. 7-amino-6-aminomethyl-5-(2,4-dichlorophenyl)-1,3-dimethyl-1H-pyrido[2,3-d]pyrimidine-2,4-dione trifluoroacetate (free base shown as II)), pharmaceutical compns., kits and methods are provided for inhibiting DPP-IV and other S9 proteases. Although the methods of preparation are not claimed, preps. and/or characterization data for .apprx.50 examples of I are included. For example, II was prepared by cyclizing 2-(2,4-dichlorobenzylidene)malononitrile (prepared from 2,4-dichlorobenzaldehyde and malononitrile) with 6-amino-1,3-dimethyluracil followed by reduction with BH₃-THF and acidification with TFA. For I: W = CR₃ and N; X = CR₄ and N; Y = CO, CS, SO, SO₂, CR₆R₆' and C:NR₆; Z = CO, CS, SO, SO₂, and C:NR₆; R₁ = (C1-10)alkyl, (C3-12)cycloalkyl, hetero(C3-12)cycloalkyl, aryl(C1-10)alkyl, heteroaryl(C1-5)alkyl, et al.; R₂ = amino(C1-6)alkyl, hetero(C3-12)cycloalkyl, hetero(C4-12)bicycloaryl, heteroaryl, and cyano; R₅ and R₇ = H, halo(C1-10)alkyl, amino, nitro, thio, sulfonamide, (C1-10)alkyl, (C3-12)cycloalkyl, et al.; addnl. details including provisos are given in the claims. Compds. I were tested according to assays for protease inhibition and observed to exhibit selective DPP-IV inhibitory activity. For example, they inhibit DPP-IV activity at concns. that are at least 50 fold less than those concns. required to produce an equiactive inhibition of protease activity for FAP α . The apparent inhibition consts. (K_i) for compds. of the invention, against DPP-IV, were .apprx.10⁻⁹ M to .apprx.10⁻⁵ M.


MSTR 1





G1 = 153 / 154 / (Specifically claimed: 137-9 138-6 / 143-9 144-6)




G2 = H / F / Cl / Br / I / alkyl <containing 1-10 C,
no H> (substd. by 3 or more G18) / NH₂ (opt. substd.) / NO₂ /
CN / SH / SO₂NH₂ / alkyl <containing 1-10 C> /
carbocycle <containing 3-12 C, non-aromatic> /
heterocycle <containing zero or more N, zero or more O,
zero or more S, 3-12 C, non-aromatic> /
alkyl <containing 1-10 C> (substd. by 1 or more aryl) /
alkyl <containing 1-5 C> (substd. by heteroaryl <containing
zero or more N, zero or more O, zero or more S>) /
aryl <containing 9-12 C, polycyclic> /
heteroaryl <containing zero or more N, zero or more O,
zero or more S, 4-12 C, polycyclic> /
alkyl <containing 1-3 C> (substd. by G22) / 84 /
alkylamino <containing 1-10 C> /
alkyl <containing 1-10 C> (substd. by NH₂ (opt. substd.)) /
aryl / heteroaryl <containing zero or more N,
zero or more O, zero or more S> / OH / alkoxy / aryloxy /
heteroaryloxy <containing zero or more N, zero or more O,
zero or more S> / 68 / 70 / 72 / 75




G3 = H / R
G4 = alkyl <containing 1-10 C> /
carbocycle <containing 3-12 C, non-aromatic> /
heterocycle <containing zero or more N, zero or more O,
zero or more S, 3-12 C, non-aromatic> /
alkyl <containing 1-10 C> (substd. by 1 or more aryl) /
alkyl <containing 1-5 C> (substd. by heteroaryl <containing
zero or more N, zero or more O, zero or more S>) /
aryl <containing 9-12 C, polycyclic> /
heteroaryl <containing zero or more N, zero or more O,
zero or more S, 4-12 C, polycyclic> /
alkyl <containing 1-3 C> (substd. by G22) / 24 /
NH₂ (opt. substd.) / aryl / heteroaryl <containing zero or
more N, zero or more O, zero or more S> / OH / alkoxy /
aryloxy / heteroaryloxy <containing zero or more N,
zero or more O, zero or more S> / alkenyl / alkynyl / CN /
86 / 88 / 90 / 93 / (Specifically claimed: heterocycle
<containing 1-4 heteroatoms, up to 1 O, up to 1 S,
up to 4 N (no other heteroatoms), 1-4 C,
attached through 1 or more C, aromatic, 2 double bonds,
5-membered monocyclic ring> (opt. substd.) /
pyridyl (opt. substd. by 1 or more G13) /
Ph (opt. substd. by 1 or more G16) /
thienyl (opt. substd. by 1 or more G13) / 62 / 165 / 174)




G5 = alkyl <containing 1-6 C>
 (substd. by NH2 (opt. substd.)) /
 heterocycle <containing zero or more N, zero or more O,
 zero or more S, 3-12 C, non-aromatic> /
 heteroaryl <containing zero or more N, zero or more O,
 zero or more S, 4-12 C, polycyclic> /
 heteroaryl <containing zero or more N, zero or more O,
 zero or more S> / CN / (Specifically claimed: 11)




G6 = N / 27



G7 = heterocycle <containing 2-4 heteroatoms, 2 N,
 up to 2 S (no other heteroatoms), 2 or more C,
 attached through 2 or more C, 1-2 double bonds,
 6-membered monocyclic ring> (opt. substd.)
 G8 = 40 / any ring <containing zero or more N,
 zero or more O, zero or more S, attached through 1 or more C>
 (opt. substd.)




G9 = H / F / Cl / Br / I / alkyl <containing 1-10 C,
 no H> (substd. by 3 or more G18) / CN / NO2 / alkyl /
 carbocycle <non-aromatic> / alkenyl / alkynyl / aryl /
 heteroaryl <containing zero or more N, zero or more O,
 zero or more S>
 G10 = H / R
 G11 = NH2 / 46 / 48 / heterocycle <containing 1 or more
 N, zero or more O, zero or more S, attached through 1 N>



G12 = alkyl <containing 1-10 C, no H> (substd. by 3 or more G18) / NH2 (opt. substd.) / SH / alkyl <containing 1-10 C> / carbocycle <containing 3-12 C, non-aromatic> / heterocycle <containing zero or more N, zero or more O, zero or more S, 3-12 C, non-aromatic> / alkyl <containing 1-10 C> (substd. by 1 or more aryl) / alkyl <containing 1-5 C> (substd. by heteroaryl <containing zero or more N, zero or more O, zero or more S>) / aryl <containing 9-12 C, polycyclic> / heteroaryl <containing zero or more N, zero or more O, zero or more S, 4-12 C, polycyclic> / alkyl <containing 1-3 C> (substd. by G22) / 51 / aryl / heteroaryl <containing zero or more N, zero or more O, zero or more S> / OH / alkoxy / aryloxy / heteroaryloxy <containing zero or more N, zero or more O, zero or more S>



G13 = F / Cl / Br / I / alkyl <containing 1-10 C, no H> (substd. by 3 or more G18) / CF3 / alkyl <containing 1-10 C> / alkenyl / alkynyl / aryl / heteroaryl <containing zero or more N, zero or more O, zero or more S> / SO2NH2 / alkylsulfonyl / arylsulfonyl / heteroarylsulfonyl <containing zero or more N, zero or more O, zero or more S> / aryloxy / heteroaryloxy <containing zero or more N, zero or more O, zero or more S> / alkyl (substd. by 1 or more aryl) / alkyl (substd. by heteroaryl <containing zero or more N, zero or more O, zero or more S>) / carbocycle <non-aromatic> / heterocycle <containing zero or more N, zero or more O, zero or more S, non-aromatic> / NH2 (opt. substd.) / SH / CN / NO2 / OH / alkoxy / 122 / 124 / 126 / 129




G14 = N / 54 / (Specifically claimed: 56)




G16 = F / Cl / Br / I / alkyl <containing 1-10 C, no H> (substd. by 3 or more G18) / CF3 /


alkyl <containing 1-10 C> / alkenyl / alkynyl / aryl /
heteroaryl <containing zero or more N, zero or more O,
zero or more S> / SO₂NH₂ / alkylsulfonyl / arylsulfonyl /
heteroarylsulfonyl <containing zero or more N,
zero or more O, zero or more S> / aryloxy /
heteroaryloxy <containing zero or more N, zero or more O,
zero or more S> / alkyl (substd. by 1 or more aryl) /
alkyl (substd. by heteroaryl <containing zero or more N,
zero or more O, zero or more S>) /
carbocycle <non-aromatic> / heterocycle <containing zero or
more N, zero or more O, zero or more S, non-aromatic> /
NH₂ (opt. substd.) / SH / CN / NO₂ / OH / alkoxy / 113 /
115 / 117 / 120 / OMe / CO₂H / CONH₂ / tetrazolyl / CH₂NH₂ /
CH₂OH / Ph / tetrazolyl / alkylaminocarbonyl <containing 1-7
C> / alkoxycarbonyl <containing 1-7 C>




G17 = O / S / NH (opt. substd.)
G18 = F / Cl / Br / I
G19 = H / R
G20 = S(O) / SO₂
G21 = O / S
G22 = 20 / 77 / 79 / 82



G23 = carbon chain <containing 1-3 C, saturated>
(opt. substd.)
G24 = NH (opt. substd.)
G25 = O / S / NH (opt. substd.)
G26 = H / alkyl <containing 1-10 C>
(substd. by 1 or more G18) / NH₂ (opt. substd.) / NO₂ / SH /
SO₂NH₂ / alkyl <containing 1-10 C> /
carbocycle <containing 3-12 C, non-aromatic> /
heterocycle <containing zero or more N, zero or more O,
zero or more S, 3-12 C, non-aromatic> /
alkyl <containing 1-10 C> (substd. by 1 or more aryl) /
alkyl <containing 1-5 C> (substd. by heteroaryl <containing
zero or more N, zero or more O, zero or more S>) /
aryl <containing 9-12 C, polycyclic> /
heteroaryl <containing zero or more N, zero or more O,
zero or more S, 4-12 C, polycyclic> /
alkyl <containing 1-3 C> (substd. by G22) / 197 / aryl /
heteroaryl <containing zero or more N, zero or more O,
zero or more S> / OH / alkoxy / aryloxy /
heteroaryloxy <containing zero or more N, zero or more O,
zero or more S> / 104 / 106 / 108 / 111 /
(Specifically claimed: Me)



G27 = NH₂ / 192 / 194 / heterocycle <containing 1 or more N, attached through 1 N>



G28 = NH₂ (opt. substd.) / alkyl / carbocycle <non-aromatic> / heterocycle <containing zero or more N, zero or more O, zero or more S> / alkyl (substd. by 1 or more aryl) / alkyl (substd. by heteroaryl <containing zero or more N, zero or more O, zero or more S>) / aryl <polycyclic> / heteroaryl <containing zero or more N, zero or more O, zero or more S, polycyclic>


G29 = H / R

G30 = H / R

G31 = H / R

G32 = H / R

G33 = H / F / Cl / Br / I / alkyl <containing 1-10 C, no H> (substd. by 3 or more G18) / NH₂ (opt. substd.) / NO₂ / CN / SH / SO₂NH₂ / alkyl <containing 1-10 C> / carbocycle <containing 3-12 C, non-aromatic> / heterocycle <containing zero or more N, zero or more O, zero or more S, 3-12 C, non-aromatic> / alkyl <containing 1-10 C> (substd. by 1 or more aryl) / alkyl <containing 1-5 C> (substd. by heteroaryl <containing zero or more N, zero or more O, zero or more S>) / aryl <containing 9-12 C, polycyclic> / heteroaryl <containing zero or more N, zero or more O, zero or more S, 4-12 C, polycyclic> / alkyl <containing 1-3 C> (substd. by G22) / 199 / alkylamino <containing 1-10 C> / alkyl <containing 1-10 C> (substd. by NH₂ (opt. substd.)) / aryl / heteroaryl <containing zero or more N, zero or more O, zero or more S> / OH / alkoxy / aryloxy / heteroaryloxy <containing zero or more N, zero or more O, zero or more S> / 95 / 97 / 99 / 102 / 131



G29+G30= bond

G31+G32= bond

Patent location:

Note:

claim 1

additional substitution and ring formation also claimed

L86 ANSWER 7 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 142:455888 MARPAT
 TITLE: Halogenated coordination compounds preparation and use thereof
 INVENTOR(S): Schwaiger, Jochen; Bach, Ingrid; Stoessel, Philipp
 PATENT ASSIGNEE(S): Covion Organic Semiconductors G.m.b.H., Germany
 SOURCE: PCT Int. Appl., 23 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005042545	A2	20050512	WO 2004-EP11891	20041021
WO 2005042545	A3	20060105		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				


DE 10351556 A1 20050602 DE 2003-10351556 20031103

PRIORITY APPLN. INFO.: DE 2003-10351556 20031103

OTHER SOURCE(S): CASREACT 142:455888


AB The invention relates to novel halogenated coordination compds. containing azanaphthalenols/thiols/selenols, of application as functional materials in differing applications within the widest sense of the electronic industry and which comprise organically-bonded halides as reactive groups. The invention further relates to a method for regioselective preparation of said compds. For example, All2 (HL = 8-hydroxyquinoline) was brominated with N-bromosuccinimide to give All13 (HL1 = 5-bromo-8-hydroxyquinoline) in 99 % yield.

MSTR 1





G1 = C1 / F / Br / I / H


G2 = N / 12



G3 = H / F / Cl / Br / I / CN / CHO / 16 / 22 / 24 /
 NO2 / 27 / 30 / 33 / carbon chain <containing 1 or more C>
 (opt. substd.) / 36



G4 = OH / SH / 14 / aryl <containing up to 40 C>
 (opt. substd.) / heteroaryl <containing up to 40 atoms,
 zero or more O, zero or more S,
 zero or more N (no other heteroatoms)> (opt. substd.)



G5 = H / R

G6 = carbon chain <containing 1 or more C>
 (opt. substd.) / aryl <containing up to 40 C>
 (opt. substd.) / heteroaryl <containing up to 40 atoms,
 zero or more O, zero or more S,
 zero or more N (no other heteroatoms)> (opt. substd.)

G10 = R <"metal"> / (Specifically claimed: Al / Be / B /
 Ca / Ga / In / Ir / Li / Mg / Zn)

Patent location: claim 1

Note: as complexes with G10

Note: additional ligands and ring formation also claimed

Note: also incorporates claim 13, structure IV

Maybe L86 ANSWER 8 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 142:134921 MARPAT

TITLE: Process for synthesizing quinazoline
 γ -methylene-L-glutamic acid and related
 antifolates

INVENTOR(S): Wu, Ye; Kochat, Harry

PATENT ASSIGNEE(S): Bionumerik Pharmaceuticals, Inc., USA

SOURCE: U.S. Pat. Appl. Publ., 8 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2005020833	A1	20050127	US 2003-627483	20030725
US 7060825	B2	20060613		

PRIORITY APPLN. INFO.: US 2003-627483 20030725


GI




AB The invention relates to a process for synthesizing antifolate compds. I [R1, R3 are individually NH2, alkylamino, OH, alkoxy, keto, alkyl or a protecting group; R3 is H, OH, alkoxy, trifluoromethylalkoxy, halo or alkylthio; R4 is OH, alkoxy or CO-X, where X is OH, alkoxy or an amino acid residue; X1-X4 are each individually carbon or nitrogen] that have com. use as drugs in oncol., inflammatory disease, and other medical fields. Thus, cyclization reaction of 2-amino-5-nitrobenzonitrile with guanidine in the presence of NaOMe afforded 2,4-diamino-6-nitroquinazoline (Q-NO2). The latter is converted to reactive aldehyde Q-CHO in 3 steps starting with reduction to the amine. The 6-formylquinazoline is then coupled to a benzoic acid ester by a modified Horner reaction. Subsequent hydrogenation, saponification, coupling with di-Et γ -methylene-L-glutamate, and saponification afforded p-Q-CH₂CH₂C₆H₄CONHCH(CO₂H)CH₂C(:CH₂)CO₂H- (S).

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1




G1 = 18 / 31



G2 = CH / N

G3 = NH2 / alkylamino / dialkylamino / OH / alkoxy / 37
/ loweralkyl / R <"nitrogen or oxygen protecting group">



G4 = H / OH / alkoxy / CF3 / halo / SH / alkylthio



G5 = H / alkoxy / 40

US 2003130278	A1	20030710	US 2002-269197	20021011
US 6962922	B2	20051108		
BR 2002013239	A	20040928	BR 2002-13239	20021011
EP 1465878	A1	20041013	EP 2002-801341	20021011
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2005509626	T2	20050414	JP 2003-536218	20021011
US 2005245548	A1	20051103	US 2005-148880	20050609

PRIORITY APPLN. INFO.:

US 2001-329181P	20011012
WO 2001-EP11824	20011012
US 2002-395441P	20020712
WO 2002-EP8475	20020712
US 2002-269197	20021011
WO 2002-EP12194	20021011


GI




AB The title compds. I [W1 = O, S, (substituted)amino; W2 = H, CF3, (substituted)amino, alkyl, alkenyl, alkynyl, aryl, etc.; W1W2 = heteroalkylene, etc; X1, X2 and X3 = N or (substituted)carbon; n = 0-8; Z = CR3R4, where R3, R4 = H, alkyl, halogen, (substituted)amino, etc.; A = (hetero)aryl or (hetero)cycloalkyl; R1 = H, alkyl, alkenyl, alkynyl, etc.; R2 = H, alkyl, halogen, cyano, nitro, (substituted)amino, etc.; q = 0-7] were prepared as inhibitors of type-13 matrix metalloprotease. Thus, reaction of 3-(3,4-difluoro-benzyl)-6-iodo-1-methyl-1H-quinazoline-2,4-dione (preparation given) with 3-phenyl-propyne yielded compound II. The IC50 values on MMP-13 of the prepared compds. are all below 10 μ M.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1



G6 = OH / alkoxy / R <"amino acid residue"> /
(Example: 42)




G7 = (1-2) CH₂


Patent location: claim 1

1 L86 ANSWER 9 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 140:94063 MARPAT
 TITLE: Preparation of new alkynylated quinazoline compounds
 as MMP-13 inhibitors
 INVENTOR(S): Gaudilliere, Bernard; Jacobelli, Henry; Wilson,
 Michael William; Picard, Joseph Armand
 PATENT ASSIGNEE(S): Warner-Lambert Company Llc, USA
 SOURCE: PCT Int. Appl., 77 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007469	A1	20040122	WO 2002-EP8475	20020712
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002331362	A1	20040202	AU 2002-331362	20020712
CA 2463159	AA	20030424	CA 2002-2463159	20021011
WO 2003033478	A1	20030424	WO 2002-EP12194	20021011
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				




G1 = 1-19 9-21 / 104-19 112-21



G2 = alkyl <containing 1-6 C> / OH / CN

G3 = H / CF₃ / NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> /
alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4) /
alkyl <containing 1-6 C> (substd. by 1 or more G30) /
alkyl <containing 1-6 C> (substd. by G31) /
heteroaryl <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
monocyclic> (opt. substd. by 1 or more G4) /
heterocycle <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G4) / (Specifically claimed: Me)


G4 = halo / NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / CN /
alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
SH / alkoxy <containing 1-6 C> /
alkylthio <containing 1-6 C>




G5 = alkyl <containing 1-6 C> / Ph / OH /

alkoxy <containing 1-16 C>

G6 = O / S / NH / 13




G7 = (up to 2) N / 22




G8 = H / alkyl <containing 1-6 C> / OH /

alkoxy <containing 1-6 C> / halo / CF₃ / CN / NO₂ / SH / 24


/
 26 / NH2 / 28 / 30 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>




G9 = S / S(O)
 G10 = S / S(O) / SO2
 G11 = alkyl <containing 1-6 C>
 G12 = alkyl <containing 1-6 C> /
 alkyl <containing 1-6 C> (substd. by 1 or more aryl
 <containing up to 10 C, mono- or bicyclic>)
 G13 = ethynylene / 33-18 35-20





G14 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 36 / 38-18 40-34 /
 (Specifically claimed: CH2)






G15 = halo / NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
 CO2H / alkoxycarbonyl <containing 1-6 C>
 G16 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G17 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G18 = O / S / S(O) / SO2 / NH / 41




G19 = alkyl <containing 1-6 C>
 G20 = carbocycle <containing 8-10 C,
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 (Specifically claimed: Ph / imidazolyl / 239 / 243 / 247 /
 254 / 258 / 266 / 273)




G21 = alkyl <containing 1-6 C> / halo / CN / NO2 /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 43 /
 45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
 73 / 76 / CO2H / alkoxycarbonyl <containing 1-6 C> / 80 /
 83 / 86 / 90 / 92 / 95 / aryl <containing up to 10 C,
 mono- or bicyclic> (opt. substd. by (1-4) G29) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 mono- or bicyclic> (opt. substd. by (1-4) G29) /
 heterocycle <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds, mono- or bicyclic,
 (1) 5- or more membered ring, (1) up to 6-membered ring>
 (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,
 non-aromatic, 0 or more double bonds, mono- or bicyclic>
 (opt. substd. by 1 or more G29) / 168






- G22 = alkyl <containing 1-6 C> / Ph
 G23 = (1-3) CH2
 G24 = NH2 / 56 / 58 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring>



- G25 = O / S / NH / 66



- G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH2 / O / S / S(O) / SO2
 G28 = aryl <containing up to 10 C, mono- or bicyclic> (opt. substd. by (1-4) G29) / heteroaryl <containing 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), mono- or bicyclic> (opt. substd. by (1-4) G29) / heterocycle <containing 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), non-aromatic, 0 or more double bonds, mono- or bicyclic, (1) 5- or more membered ring, (1) up to 6-membered ring> (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C, non-aromatic, 0 or more double bonds, mono- or bicyclic> (opt. substd. by 1 or more G29)
 G29 = alkyl <containing 1-6 C> / halo / alkyl <containing 1-6 C> (substd. by (3) halo) / OH / alkoxy <containing 1-6 C> / SH / alkylthio <containing 1-6 C> / NH2 / alkylamino <containing 1-6 C> / dialkylamino <each alkyl containing 1-6 C>
 G30 = aryl <containing up to 10 C, mono- or bicyclic> (opt. substd. by 1 or more G4)
 G31 = carbocycle <containing 3-10 C, non-aromatic, 0 or more double bonds, mono- or bicyclic> (opt. substd. by 1 or more G4)
 G32 = H / alkyl <containing 1-6 C> (opt. substd. by 1 or more G33) / alkenyl <containing 2-6 C> (opt. substd. by 1 or more G33) / alkynyl <containing 2-6 C> (opt. substd. by 1 or more G33) / 97 / carbocycle <containing 5-10 C, 0 or more double bonds, mono- or bicyclic, 5- or 6-membered rings only> (opt. substd. by (1-7) G41) / heterocycle <containing 5-10 atoms, 1-4 heteroatoms, zero or more N, zero or more O,

zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41)

^{G34-G37}
_{97 98}





G33 = NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
 CO₂H / alkoxycarbonyl <containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 99 / 101-20 103-98 /
 (Specifically claimed: CH₂)

^{G16=O} ^{G17-G18-G17}
_{99 101 103}

G35 = N / 118






^{C-G36}
₁₁₈

G36 = H / R
 G37 = carbocycle <containing 5-10 C,
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
 (Specifically claimed: 285 / 294 / 301 / 308)




G38 = (0-3) CH₂
 G40 = alkyl <containing 1-6 C>
 G41 = alkyl <containing 1-6 C> / halo / CN / NO₂ /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH₂ / 232 /
 234 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / OH / SH / 173 / 175 / 177 / 179 / 181 / 183 / 186 /
 189 / 191 / 194 / CO₂H / alkoxycarbonyl <containing 1-6 C> /
 198 / 201 / 204 / 208 / 210 / 213 / 215 /
 aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by (1-4) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>


(opt. substd. by (1-4) G29) / heterocycle <containing 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), non-aromatic, 0 or more double bonds, mono- or bicyclic, (1) 5- or more membered ring, (1) up to 6-membered ring>
 (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C, non-aromatic, 0 or more double bonds, mono- or bicyclic>
 (opt. substd. by 1 or more G29)




G42 = alkylene <containing 1-6 C>
 G43 = OH / alkoxy <containing 1-6 C> / NH2 / 220 / 222 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring> / 225




G44 = OH / alkoxy <containing 1-6 C> / NH2 / 227 / 229 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring>




G45 = OMe / F / 278 / Ph / Cl



G46 = F / Cl / 290



G47 = OMe / OBu-t / OH / 316 / NMe2 / piperidino / NHet /
324




G48 = Cl / F
Patent location:
Note:

claim 1
and N-oxides or pharmaceutically acceptable acid or
base addition salts


Note: additional heteroatom interruptions in G17 also
claimed

Note: substitution is restricted
Stereochemistry: and optical isomers

MSTR 2



G1 = 1-19 9-21 / 104-19 112-21



G2 = alkyl <containing 1-6 C> / OH / CN
G3 = H / CF3 / NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> /
alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4) /

alkyl <containing 1-6 C> (substd. by 1 or more G30) /
 alkyl <containing 1-6 C> (substd. by G31) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 monocyclic> (opt. substd. by 1 or more G4) /
 heterocycle <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G4) / (Specifically claimed: Me)
 G4 = halo / NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / CN /
 alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
 SH / alkoxy <containing 1-6 C> /
 alkylthio <containing 1-6 C>

$\overset{\text{C}}{\underset{16}{\text{O}}}$ -G5

G5 = alkyl <containing 1-6 C> / Ph / OH /
 alkoxy <containing 1-16 C>
 G6 = O / S / NH / 13

$\overset{\text{N}}{\underset{13}{\text{---}}}$ -G2

G7 = (up to 2) N / 22

$\overset{\text{C}}{\underset{22}{\text{---}}}$ -G8

G8 = H / alkyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / halo / CF₃ / CN / NO₂ / SH / 24
 /
 26 / NH₂ / 28 / 30 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>




$\overset{\text{G9}}{\underset{24}{\text{---}}}$ -OH

$\overset{\text{G10}}{\underset{26}{\text{---}}}$ -G11


$\overset{\text{HN}}{\underset{28}{\text{---}}}$ -G12

$\overset{\text{G12}}{\underset{\text{N}}{\text{---}}} \begin{matrix} \text{G12} \\ \text{30} \end{matrix}$



G9 = S / S(O)
 G10 = S / S(O) / SO₂
 G11 = alkyl <containing 1-6 C>
 G12 = alkyl <containing 1-6 C> /
 alkyl <containing 1-6 C> (substd. by 1 or more aryl
 <containing up to 10 C, mono- or bicyclic>)
 G13 = H / halo / 237 / CHO / COMe / R <"ester group"> /
 240 / ethynyl / 246




- G15 = halo / NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH / SH /
alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
CO₂H / alkoxycarbonyl <containing 1-6 C>
- G16 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
- G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
- G18 = O / S / S(O) / SO₂ / NH / 41



- G19 = alkyl <containing 1-6 C>
- G22 = alkyl <containing 1-6 C> / Ph
- G23 = (1-3) CH₂
- G24 = NH₂ / 56 / 58 / heterocycle <containing 1-2
heteroatoms, 1 or more N, zero or more O (no other
heteroatoms), attached through 1 or more N,
5- to 6-membered monocyclic ring>



- G25 = O / S / NH / 66



- G26 = OH / alkoxy <containing 1-6 C>
- G27 = CH₂ / O / S / S(O) / SO₂
- G28 = aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by (1-4) G29) / heteroaryl <containing 1-4
heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms), mono- or bicyclic>
(opt. substd. by (1-4) G29) / heterocycle <containing 1-4
heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms), non-aromatic,
0 or more double bonds, mono- or bicyclic,
(1) 5- or more membered ring, (1) up to 6-membered ring>
(opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,
non-aromatic, 0 or more double bonds, mono- or bicyclic>
(opt. substd. by 1 or more G29)
- G29 = alkyl <containing 1-6 C> / halo /
alkyl <containing 1-6 C> (substd. by (3) halo) / OH /

alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G30 = aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G4)
 G31 = carbocycle <containing 3-10 C, non-aromatic,
 0 or more double bonds, mono- or bicyclic>
 (opt. substd. by 1 or more G4)
 G32 = H / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G33) /
 alkenyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
 alkynyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
 97 / carbocycle <containing 5-10 C, 0 or more double bonds,
 mono- or bicyclic, 5- or 6-membered rings only>
 (opt. substd. by (1-7) G41) / heterocycle <containing 5-10
 atoms, 1-4 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41)

~~G34-G37~~
~~97 98~~

G33 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
 CO2H / alkoxy carbonyl <containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 99 / 101-20 103-98 /
 (Specifically claimed: CH2)

~~G16=O~~ ~~G17-G18-G17~~
~~99~~ ~~101~~ ~~103~~

G35 = N / 118

~~C-G36~~
~~118~~

G36 = H / R
 G37 = carbocycle <containing 5-10 C,
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41)
 G38 = (0-3) CH2
 G41 = alkyl <containing 1-6 C> / halo / CN / NO2 /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 232 /
 234 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic

ring> / OH / SH / 173 / 175 / 177 / 179 / 181 / 183 / 186 /
 189 / 191 / 194 / CO2H / alkoxy carbonyl <containing 1-6 C> /
 198 / 201 / 204 / 208 / 210 / 213 / 215 /
 aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by (1-4) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>
 (opt. substd. by (1-4) G29) / heterocycle <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, mono- or bicyclic,
 (1) 5- or more membered ring, (1) up to 6-membered ring>
 (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,
 non-aromatic, 0 or more double bonds, mono- or bicyclic>
 (opt. substd. by 1 or more G29)

G9—OH G10—G11 C(O)—G22 G23—G24 G25—G24
 173 175 177 179 181

G25—G23—G24 G23—SO₂—G24 O₂S—G24 G25—C(O)—G26
 183 186 189 191

G25—G23—C(O)—G26 G23—C(O)—G26 G25—C(O)—G24
 194 198 201


G25—G23—C(O)—G24 C(O)—G24 G23—C(O)—G24 G27—G28
 204 208 210 213

G38—C(O)—O—G42—G43 HN—G12
 215 232
 N
 234
 G12
 G12

G42 = alkylene <containing 1-6 C>
 G43 = OH / alkoxy <containing 1-6 C> / NH₂ / 220 / 222 /
 heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / 225

HN—G12 G12
 220 222
 N
 222
 G12 C(O)—G44
 225

G44 = OH / alkoxy <containing 1-6 C> / NH₂ / 227 / 229 /
 heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring>



G45 = Me / CF3

Patent location:

Note: claim 12
additional heteroatom interruptions in G17 also
claimed

Note: substitution is restricted

L86 ANSWER 10 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 140:105274 MARPAT

TITLE: Combination of an allosteric alkyne inhibitor of
matrix metalloproteinase-13 with a selective inhibitor
of cyclooxygenase-2

INVENTOR(S): Roark, William Howard

PATENT ASSIGNEE(S): Warner-Lambert Company Llc, USA

SOURCE: PCT Int. Appl., 196 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007025	A1	20040122	WO 2003-IB3043	20030707
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2491820	AA	20040122	CA 2003-2491820	20030707
AU 2003281168	A1	20040202	AU 2003-281168	20030707
EP 1525030	A1	20050427	EP 2003-740952	20030707
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2003012666	A	20050510	BR 2003-12666	20030707
JP 2006502991	T2	20060126	JP 2004-520997	20030707
US 2004019055	A1	20040129	US 2003-620173	20030715
PRIORITY APPLN. INFO.:			US 2002-396385P	20020717
			WO 2003-IB3043	20030707

GI




AB The invention provides a combination, comprising an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with a selective inhibitor of COX-2, or a pharmaceutically acceptable salt thereof, that is not celecoxib or valdecoxib. This invention also provides a combination comprising an NSAID, or a pharmaceutically acceptable salt thereof, and an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof. This invention also provides a method of treating a disease that is responsive to inhibition of MMP-13 and cyclooxygenase 1 or cyclooxygenase-2, comprising administering to a patient suffering from such a disease the invention combination comprising an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with an NSAID, or a pharmaceutically acceptable salt thereof. Biol. examples include methods to determine inhibition of MMP-13 and COX-2 by the compds. and chemical-induced arthritis in animals. Pharmaceutical formulations are also given. An example MMP-13 inhibitor is I.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1

G20-G13-G1-G32
18 19 20 21

G1 = 1-19 9-21 / 104-19 112-21



G2 = alkyl <containing 1-6 C> / OH / CN
G3 = H / CF3 / NH2 / alkylamino <containing 1-10 C> /
dialkylamino <each alkyl containing 1-10 C> /
alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4) /
alkyl <containing 1-10 C> (substd. by 1 or more G30) /
alkyl <containing 1-10 C> (substd. by G31) /
heteroaryl <containing 1-4 heteroatoms, zero or more N,

zero or more O, zero or more S (no other heteroatoms),
monocyclic> (opt. substd. by 1 or more G4) /
heterocycle <containing 1-3 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G4) / (Specifically claimed: Me)
G4 = halo / NH₂ / alkylamino <containing 1-10 C> /
dialkylamino <each alkyl containing 1-10 C> / CN /
alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
SH / alkoxy <containing 1-6 C> /
alkylthio <containing 1-6 C>

$\text{C}(\text{O})\text{-G5}$
₁₆

G5 = alkyl <containing 1-6 C> / Ph / OH /
alkoxy <containing 1-16 C>
G6 = O / S / NH / 13

$\text{N}\text{---G2}$
₁₃

G7 = (up to 2) N / 22

$\text{C}\text{---G8}$
₂₂

G8 = H / alkyl <containing 1-6 C> / OH /
alkoxy <containing 1-6 C> / halo / CF₃ / CN / NO₂ / SH / 24
/ 26 / NH₂ / 28 / 30 / heterocycle <containing 1-2
heteroatoms, 1 or more N, zero or more O (no other
heteroatoms), attached through 1 or more N,
5- to 6-membered monocyclic ring>
(opt. substd. by alkyl <containing 1-10 C>)

$\text{G9}\text{---OH}$
₂₄

$\text{G10}\text{---G11}$
₂₆

$\text{HN}\text{---G12}$
₂₈


$\text{N}\text{---G12}$
₃₀

G9 = S / S(O)
G10 = S / S(O) / SO₂
G11 = alkyl <containing 1-6 C>
G12 = alkyl <containing 1-6 C> /
alkyl <containing 1-6 C> (substd. by 1 or more aryl
<containing up to 10 C, mono- or bicyclic>)
G13 = ethynylene / 33-18 35-20


$\text{G14}\text{---C}\equiv\text{C}$
_{33 34 35}

G14 = carbon chain <containing 1 or more C,
up to 2 double bonds, up to 2 triple bonds>
(opt. substd. by 1 or more G15) / 36 / 38-18 40-34 /


(Specifically claimed: CH2)



- G15 = halo / NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH / SH /
alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
CO2H / alkoxycarbonyl <containing 1-6 C>
- G16 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
- G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
- G18 = O / S / S(O) / SO2 / NH / 41






- G19 = alkyl <containing 1-6 C>
- G20 = aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by (1-7) G21) / heterocycle <containing 1-4
heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G21) / carbocycle <containing 5-6 C,
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G21) / carbocycle <containing 8-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
(Specifically claimed: 239 / Ph)




- G21 = alkyl <containing 1-10 C> / halo / CN / NO2 /
alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 43 /
45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
zero or more O (no other heteroatoms),
attached through 1 or more N, 5- to 6-membered monocyclic
ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
73 / 76 / CO2H / alkoxycarbonyl <containing 1-6 C> / 80 /
83 / 86 / 90 / 92 / 95 / carbocycle <containing 6 C,
aromatic, 6 normalized bonds, 6-membered monocyclic ring>
(opt. substd. by (1-3) G29) / heteroaryl <containing 1-4
heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms), monocyclic>
(opt. substd. by (1-3) G29) / cycloalkyl <containing 5-6 C>
(opt. substd. by (1-3) G29) / heterocycle <containing 1-3


heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29)



G22 = alkyl <containing 1-6 C> / Ph
 G23 = (1-3) CH2
 G24 = NH2 / 56 / 58 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>




G25 = O / S / NH / 66



G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH2 / O / S / S(O) / SO2
 G28 = carbocycle <containing 6 C, aromatic,
 6 normalized bonds, 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), monocyclic>
 (opt. substd. by (1-3) G29) / cycloalkyl <containing 5-6 C>
 (opt. substd. by (1-3) G29) / heterocycle <containing 1-3
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29)
 G29 = alkyl <containing 1-10 C> / halo /
 alkyl <containing 1-6 C> (substd. by (3) halo) / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / NH2 /

- alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C>
G30 = aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4)
G31 = cycloalkyl <containing 3-10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4)
G32 = H / alkyl <containing 1-6 C>
(opt. substd. by 1 or more G33) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
97 / carbocycle <containing 5-10 C, 0 or more double bonds,
mono- or bicyclic, 5- or 6-membered rings only>
(opt. substd. by (1-7) G41) / heterocycle <containing 5-10
atoms, 1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
(Specifically claimed: 250 / 259 / CH2Ph)

G34-G37
97 98



- G33 = NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> /
alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
CO2H / alkoxycarbonyl <containing 1-6 C> / OH / SH /
alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
G34 = carbon chain <containing 1 or more C,
up to 2 double bonds, up to 2 triple bonds>
(opt. substd. by 1 or more G15) / 99 / 101-20 103-98

G16=O
99

G17-G18-G17
101 103

- G35 = N / 118

C-G36
118

- G36 = H / R
G37 = carbocycle <containing 5-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41)
G41 = alkyl <containing 1-6 C> / halo / CN / NO2 /
alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 232 /
234 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
zero or more O (no other heteroatoms),
attached through 1 or more N, 5- to 6-membered monocyclic
ring> / OH / SH / 173 / 175 / 177 / 179 / 181 / 183 / 186 /
189 / 191 / 194 / CO2H / alkoxycarbonyl <containing 1-6 C> /
198 / 201 / 204 / 208 / 210 / 213 /


carbocycle <containing 6 C, aromatic, 6 normalized bonds,
 6-membered monocyclic ring> (opt. substd. by (1-3) G29) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 monocyclic> (opt. substd. by (1-3) G29) /
 cycloalkyl <containing 5-6 C> (opt. substd. by (1-3) G29) /
 heterocycle <containing 1-3 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29)

$\text{G9}-\text{OH}$ $\text{G10}-\text{G11}$ $\text{C}(\text{O})-\text{G22}$ $\text{G23}-\text{G24}$ $\text{G25}-\text{G24}$
 173 175 177 179 181

$\text{G25}-\text{G23}-\text{G24}$ $\text{G23}-\text{SO}_2-\text{G24}$ $\text{O}_2\text{S}-\text{G24}$ $\text{G25}-\text{C}(\text{O})-\text{G26}$
 183 186 189 191

$\text{G25}-\text{G23}-\text{C}(\text{O})-\text{G26}$ $\text{G23}-\text{C}(\text{O})-\text{G26}$ $\text{G25}-\text{C}(\text{O})-\text{G24}$
 194 198 201

$\text{G25}-\text{G23}-\text{C}(\text{O})-\text{G24}$ $\text{C}(\text{O})-\text{G24}$ $\text{G23}-\text{C}(\text{O})-\text{G24}$ $\text{G27}-\text{G28}$
 204 208 210 213




Patent location: claim 1
 Note: or pharmaceutically acceptable salts or N-oxides
 Note: additional heteroatom interruptions in G17 also
 claimed
 Note: substitution is restricted
 Note: additional ring formation also claimed

L86 ANSWER 11 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 140:105336 MARPAT
 TITLE: Combination of an allosteric carboxylic inhibitor of
 matrix metalloproteinase-13 with a selective inhibitor
 of cyclooxygenase-2 that is not celecoxib or
 valdecoxib, and therapeutic use
 INVENTOR(S): Roark, William Howard
 PATENT ASSIGNEE(S): Warner-Lambert Company LLC, USA
 SOURCE: PCT Int. Appl., 239 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004006931	A2	20040122	WO 2003-IB3098	20030707
WO 2004006931	A3	20040513		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2492387	AA	20040122	CA 2003-2492387	20030707
AU 2003281170	A1	20040202	AU 2003-281170	20030707
BR 2003012744	A	20050426	BR 2003-12744	20030707
EP 1530475	A2	20050518	EP 2003-740981	20030707
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2006503812	T2	20060202	JP 2004-521004	20030707
US 2004019054	A1	20040129	US 2003-619769	20030715
PRIORITY APPLN. INFO.:			US 2002-396785P	20020717
			WO 2003-IB3098	20030707

AB The invention provides a combination comprising an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with a selective inhibitor of COX-2, or a pharmaceutically acceptable salt thereof, that is not celecoxib or valdecoxib, and their use for the treatment of diseases that are responsive to inhibition of MMP-13 and cyclooxygenase-2.

MSTR 1



- G1 = H / NH₂ / alkyl <containing 1-6 C> (opt. substd.) / alkenyl <containing 3-6 C> (opt. substd.) / alkynyl <containing 3-6 C> (opt. substd.) / alkyl <containing 1-6 C> (substd. by alkylamino <containing 1-6 C> (opt. substd.)) / alkyl <containing 1-6 C> (substd. by dialkylamino <each alkyl containing 1-6 C> (opt. substd.)) / aryl <containing up to 10 C, mono- or bicyclic> (opt. substd.) / alkyl <containing 1-6 C> (substd. by 1 or more G2) / heterocycle <containing 5-6 atoms, 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), 0 or more double bonds, 5- to 6-membered monocyclic ring> (opt. substd.) / alkyl <containing 1-6 C> (substd. by cycloalkyl <containing 3-6 C> (opt. substd.)) / (Specifically claimed: Me)
- G2 = aryl <containing up to 10 C, mono- or bicyclic>

(opt. substd.)
G3 = O / S / 14

$\text{N} \text{---} \text{G4}$
14

G4 = alkyl <containing 1-6 C> (opt. substd.) / OH / CN
G5 = (up to 2) N / 16

$\text{C} \text{---} \text{G6}$
16

G6 = H / alkyl <containing 1-6 C> (opt. substd.) / NH2 /
alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH /
alkoxy <containing 1-6 C> (opt. substd.) / F / Cl / Br / I
G7 = bond / 26

$\text{C} \text{= G8}$
26

G8 = O / S / NH / 20

$\text{N} \text{---} \text{G9}$
20

G9 = alkyl <containing 1-6 C> (opt. substd.)
G10 = 24-35 25-18 / carbon chain <containing 1 or more
C, 0 or more double bonds> (opt. substd. by 1 or more G14)

$\text{G11} \text{---} \text{G12}$
24 25

G11 = carbon chain <containing 1 or more C,
0 or more double bonds> (opt. substd. by 1 or more G14) /
(Specifically claimed: CH2)
G12 = O / S / NH / 28




$\text{N} \text{---} \text{G13}$
28

G13 = alkyl <containing 1-6 C>
(opt. substd. by 1 or more G2) /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd.) / cycloalkyl / heteroaryl <containing zero or
more N, zero or more O, zero or more S (no other heteroatoms)
> (opt. substd.)
G14 = F / Cl / Br / I / NH2 / OH / SH / 30 / 32 /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd.) / heterocycle <containing zero or more N,
zero or more O, zero or more S, 0 or more double bonds>
(opt. substd.) / cycloalkyl

G15-G16
30

C(O)-G17
32

G15 = O / S
G16 = alkyl <containing 1-6 C> (opt. substd.)
G17 = OH / alkoxy <containing 1-6 C> (opt. substd.)
G18 = carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G19) / heterocycle <containing 5-6
atoms, 1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G19) / carbocycle <containing 8-10 C,
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G19) /
heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G19) /
(Specifically claimed: 163 / 174 / 179)



G19 = alkyl <containing 1-6 C> (opt. substd.) / F / Cl /
Br / I / CN / NO2 / SCF3 / CF3 / OCF3 / NH2 /
alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
OH / SH / 36 / 38 / 40 / 42 / 45 / 48 / 54 / 56 / 59

G20-OH G21-G22 O₂S-G23 G24-SO₂-G23 G25-C(O)-G26
36 38 40 42 45

G25-G24-C(O)-G26 C(O)-G26 G24-C(O)-G26 G27-G28
48 54 56 59

G20 = S / S(O)
G21 = O / S / S(O) / SO₂
G22 = alkyl <containing 1-6 C> (opt. substd.)
G23 = NH₂ / alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.)
G24 = (1-3) CH₂
G25 = O / S / S(O) / SO₂ / NH / 52


^N—G22
52

- G26 = OH / alkoxy <containing 1-6 C> (opt. substd.) /
NH2 / alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.)
G27 = CH2 / O / S / S(O) / SO2 / NH / 61

^N—G22
61

- G28 = carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G29) /
heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G29)
G29 = alkyl <containing 1-6 C> (opt. substd.) / F / Cl /
Br / I / OH / NH2
G30 = H / carbon chain <containing 1-6 C>
(opt. substd. by 1 or more G31) / 67 /
carbocycle <containing 5-6 C, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G39) / heterocycle <containing 5-6
atoms, 1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G39) / carbocycle <containing 8-10 C,
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G39) /
heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G39) /
(Specifically claimed: 184)

^{G36-G38}
67



- G31 = NH2 / CN / alkyl <containing 1-6 C>
(substd. by 1 or more G32) / cycloalkyl / 63 / OH / SH / 65

^{C(O)}G33
63





^{G34-G35}
65



- G32 = F / Cl / Br / I
G33 = NH2 / alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
OH / alkoxy <containing 1-6 C> (opt. substd.)
G34 = O / S
G35 = alkyl <containing 1-6 C> (opt. substd.)

G36 = carbon chain <containing 1 or more C,
0 or more double bonds> (opt. substd. by 1 or more G37)
G37 = Ph / F / Cl / Br / I / NH₂ / OH / SH / 69 / 71


^{G34-G35}₆₉ ^{C(O)-G17}₇₁

G38 = carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G39) / heterocycle <containing 5-6
atoms, 1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by (1-7) G39) / carbocycle <containing 8-10 C,
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G39) /
heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G39)
G39 = alkyl <containing 1-6 C> (opt. substd.) / F / Cl /
Br / I / CN / NO₂ / CF₃ / OCF₃ /
alkylamino <containing 1-6 C> (opt. substd.) /
dialkylamino <each alkyl containing 1-6 C> (opt. substd.) /
73 / 77 / 80 / 83 / 86 / 92 / OH / SH / 102 / 104 / 106 /
110 / 112 / 115 / 118 / 123 / 126 / 129 / 133 / 138 / 141 /
146 / 149 / 152 / 155 / carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.) / heterocycle <containing 5-6 atoms,
1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.) / 157






G40 = NH / 75



G41 = H / alkyl <containing 1-6 C> (opt. substd.)


G42 = 96 / 99



G43 = (1-4) CH₂

G44 = cycloalkylene <containing 3-6 C>

G45 = carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.) / heterocycle <containing 5-6 atoms,
1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.) / 159



G47 = carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.) / heterocycle <containing 5-6 atoms,
1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd.)

G48 = CO₂H / H / CO₂Me / CN

Patent location: claim 1

Note: or pharmaceutically acceptable salts or N-oxides

Note: substitution is restricted

L86 ANSWER 12 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 140:105335 MARPAT

TITLE: Combination of an allosteric alkyne inhibitor of
matrix metalloproteinase-13 with celecoxib or
valdecoxib, pharmaceutical compositions, and
therapeutic use

INVENTOR(S): Roark, William Howard

PATENT ASSIGNEE(S): Warner-Lambert Company LLC, USA

SOURCE: PCT Int. Appl., 188 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004006914	A1	20040122	WO 2003-IB3154	20030707
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2489722	AA	20040122	CA 2003-2489722	20030707
AU 2003249505	A1	20040202	AU 2003-249505	20030707
BR 2003012708	A	20050426	BR 2003-12708	20030707
EP 1534274	A1	20050601	EP 2003-764068	20030707
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2006502114	T2	20060119	JP 2004-521017	20030707
US 2004023969	A1	20040205	US 2003-619777	20030715
PRIORITY APPLN. INFO.:			US 2002-396922P	20020717
			WO 2003-IB3154	20030707


AB The invention provides a combination comprising an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof. This invention also provides a method of treating a disease that is responsive to inhibition of MMP-13 and cyclooxygenase-2, comprising administering to a patient suffering from such a disease the invention combination comprising an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof. This invention also provides a pharmaceutical composition, comprising the invention combination comprising an allosteric alkyne inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent, or excipient. The invention combinations may also be further combined with other pharmaceutical agents depending on the disease being treated.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


MSTR 1

G20-G13-G1-G32
 18 19 20 21


G1 = 1-19 9-21 / 104-19 112-21




- G2 = alkyl <containing 1-6 C> / OH / CN
 G3 = H / CF3 / NH2 / alkylamino <containing 1-10 C> /
 dialkylamino <each alkyl containing 1-10 C> /
 alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
 alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
 alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
 aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G4) /
 alkyl <containing 1-10 C> (substd. by 1 or more G30) /
 alkyl <containing 1-10 C> (substd. by G31) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 monocyclic> (opt. substd. by 1 or more G4) /
 heterocycle <containing 1-3 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G4) / (Specifically claimed: Me)
 G4 = halo / NH2 / alkylamino <containing 1-10 C> /
 dialkylamino <each alkyl containing 1-10 C> / CN /
 alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
 SH / alkoxy <containing 1-6 C> /
 alkylthio <containing 1-6 C>




- G5 = alkyl <containing 1-6 C> / Ph / OH /
 alkoxy <containing 1-16 C>
 G6 = O / S / NH / 13




- G7 = (up to 2) N / 22




- G8 = H / alkyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / halo / CF3 / CN / NO2 / SH / 24
 /
 26 / NH2 / 28 / 30 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>
 (opt. substd. by alkyl <containing 1-10 C>)




- G9 = S / S(O)
 G10 = S / S(O) / SO2
 G11 = alkyl <containing 1-6 C>
 G12 = alkyl <containing 1-6 C> /
 alkyl <containing 1-6 C> (substd. by 1 or more aryl
 <containing up to 10 C, mono- or bicyclic>)
 G13 = ethynylene / 33-18 35-20




- G14 = carbon chain <containing 1 or more C,
 up to 2 double bonds, up to 2 triple bonds>
 (opt. substd. by 1 or more G15) / 36 / 38-18 40-34 /
 (Specifically claimed: CH2)






- G15 = halo / NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
 CO2H / alkoxycarbonyl <containing 1-6 C>
 G16 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G17 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G18 = O / S / S(O) / SO2 / NH / 41




- G19 = alkyl <containing 1-6 C>
 G20 = aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by (1-7) G21) / heterocycle <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-7) G21) / carbocycle <containing 5-6 C,
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-7) G21) / carbocycle <containing 8-10 C,
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 (Specifically claimed: 239 / Ph)



G21 = alkyl <containing 1-10 C> / halo / CN / NO2 /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 43 /
 45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
 73 / 76 / CO2H / alkoxy carbonyl <containing 1-6 C> / 80 /
 83 / 86 / 90 / 92 / 95 / carbocycle <containing 6 C,
 aromatic, 6 normalized bonds, 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), monocyclic>
 (opt. substd. by (1-3) G29) / cycloalkyl <containing 5-6 C>
 (opt. substd. by (1-3) G29) / heterocycle <containing 1-3
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29)



G22 = alkyl <containing 1-6 C> / Ph
 G23 = (1-3) CH2
 G24 = NH2 / 56 / 58 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>





G25 = O / S / NH / 66

N—G19
66

- G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH2 / O / S / S(O) / SO2
 G28 = carbocycle <containing 6 C, aromatic,
 6 normalized bonds, 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), monocyclic>
 (opt. substd. by (1-3) G29) / cycloalkyl <containing 5-6 C>
 (opt. substd. by (1-3) G29) / heterocycle <containing 1-3
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by (1-3) G29)
 G29 = alkyl <containing 1-10 C> / halo /
 alkyl <containing 1-6 C> (substd. by (3) halo) / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G30 = aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G4)
 G31 = cycloalkyl <containing 3-10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G4)
 G32 = H / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G33) /
 alkenyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
 alkynyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
 97 / carbocycle <containing 5-10 C, 0 or more double bonds,
 mono- or bicyclic, 5- or 6-membered rings only>
 (opt. substd. by (1-7) G41) / heterocycle <containing 5-10
 atoms, 1-4 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
 (Specifically claimed: 250 / 259 / CH2Ph)

G34-G37
97 98



- G33 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
 CO2H / alkoxycarbonyl <containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 up to 2 double bonds, up to 2 triple bonds>
 (opt. substd. by 1 or more G15) / 99 / 101-20 103-98

G16=O
99

G17-G18-G17
101 103

G35 = N / 118

C—G36
118

G36 = H / R
G37 = carbocycle <containing 5-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41)
G41 = alkyl <containing 1-6 C> / halo / CN / NO2 /
alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 232 /
234 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
zero or more O (no other heteroatoms),
attached through 1 or more N, 5- to 6-membered monocyclic
ring> / OH / SH / 173 / 175 / 177 / 179 / 181 / 183 / 186 /
189 / 191 / 194 / CO2H / alkoxy carbonyl <containing 1-6 C> /
198 / 201 / 204 / 208 / 210 / 213 /
carbocycle <containing 6 C, aromatic, 6 normalized bonds,
6-membered monocyclic ring> (opt. substd. by (1-3) G29) /
heteroaryl <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
monocyclic> (opt. substd. by (1-3) G29) /
cycloalkyl <containing 5-6 C> (opt. substd. by (1-3) G29) /
heterocycle <containing 1-3 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by (1-3) G29)

G9—OH G10—G11 C(O)—G22 G23—G24 G25—G24
173 175 177 179 181

G25—G23—G24 G23—SO₂—G24 O₂S—G24 G25—C(O)—G26
183 186 189 191

G25—G23—C(O)—G26 G23—C(O)—G26 G25—C(O)—G24
194 198 201

G25—G23—C(O)—G24 C(O)—G24 G23—C(O)—G24 G27—G28
204 208 210 213

HN—G12
232
 N
 / \
234 G12 G12

Patent location: claim 1

Note: or pharmaceutically acceptable salts or N-oxides
 Note: additional heteroatom interruptions in G17 also
 claimed
 Note: substitution is restricted
 Note: additional ring formation also claimed

L86 ANSWER 13 OF 26 MARPAT COPYRIGHT 2006 ACS on STN


ACCESSION NUMBER: 140:105333 MARPAT
 TITLE: Combination of an allosteric carboxylic inhibitor of
 matrix metalloproteinase-13 with celecoxib or
 valdecoxib, pharmaceutical compositions, and
 therapeutic use
 INVENTOR(S): Roark, William Howard
 PATENT ASSIGNEE(S): Warner-Lambert Company LLC, USA
 SOURCE: PCT Int. Appl., 238 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004006912	A2	20040122	WO 2003-IB3044	20030707
WO 2004006912	A3	20040603		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG CA 2495432 AA 20040122 CA 2003-2495432 20030707 AU 2003281169 A1 20040202 AU 2003-281169 20030707 BR 2003012736 A 20050426 BR 2003-12736 20030707 EP 1530467 A2 20050518 EP 2003-740953 20030707 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK JP 2006503811 T2 20060202 JP 2004-520998 20030707 US 2004019053 A1 20040129 US 2003-619662 20030715 PRIORITY APPLN. INFO.: US 2002-396903P 20020717 WO 2003-IB3044 20030707				


AB The invention provides a combination comprising an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof. The invention also provides a method of treating a disease that is responsive to inhibition of MMP-13 and cyclooxygenase 2, comprising administering to a patient suffering from such a disease the invention combination comprising an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof. The invention also provides a pharmaceutical composition, comprising the invention combination comprising an allosteric carboxylic inhibitor of MMP-13, or a pharmaceutically acceptable salt thereof, with celecoxib, or a pharmaceutically acceptable salt thereof, or valdecoxib, or a pharmaceutically acceptable salt thereof.

pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent, or excipient. The invention combination may also be further combined with other pharmaceutical agents depending on the disease being treated.


MSTR 1



- G1 = H / NH₂ / alkyl <containing 1-6 C>
(opt. substd. by 1 or more G2) /
alkenyl <containing 3-6 C> (opt. substd. by 1 or more G3) /
alkynyl <containing 3-6 C> (opt. substd. by 1 or more G3) /
aryl (opt. substd. by 1 or more G3) /
heterocycle <containing zero or more N, zero or more O,
zero or more S> (opt. substd. by 1 or more G3) /
(Specifically claimed: Me)
- G2 = alkylamino <containing 1-6 C>
(opt. substd. by 1 or more G3) /
dialkylamino <each alkyl containing 1-6 C>
(opt. substd. by 1 or more G3) /
aryl (opt. substd. by 1 or more G3) /
cycloalkyl <containing 3-6 C> (opt. substd. by 1 or more G3)
/ NH₂ / alkyl <containing 1-6 C>
(opt. substd. by 1 or more G4) / CN / CO₂H /
alkoxycarbonyl <containing 1-6 C> / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C>
- G3 = NH₂ / alkyl <containing 1-6 C>
(opt. substd. by 1 or more G4) / CN / CO₂H /
alkoxycarbonyl <containing 1-6 C> / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C>
- G4 = F / Cl / Br / I
- G5 = O / S / 16



- G6 = alkyl <containing 1-6 C> / OH / CN
- G7 = (up to 2) N / 18



- G8 = H / alkyl <containing 1-6 C> / NH₂ /
alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH /
alkoxy <containing 1-6 C> / F / Cl / Br / I
- G9 = O / S / NH / 20

$\text{N} \text{---} \text{G10}$
20

G10 = alkyl <containing 1-6 C>
G11 = O / S / NH / 24 / 26

$\text{N} \text{---} \text{G12}$ $\text{HC} \text{---} \text{G13}$
24 26

G12 = alkyl <containing 1-6 C>
(opt. substd. by 1 or more aryl) / cycloalkyl / aryl /
heteroaryl <containing zero or more N, zero or more O,
zero or more S>
G13 = H / alkyl <containing 1-6 C>
(opt. substd. by 1 or more aryl) / aryl /
heterocycle <containing zero or more N, zero or more O,
zero or more S> / cycloalkyl
G14 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by 1 or more G15) / 30-28 32-29 /
(Specifically claimed: CH2)

$\text{G17} \text{---} \text{G18} \text{---} \text{G17}$
30 32


G15 = F / Cl / Br / I / NH2 / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C> / CO2H /
alkoxycarbonyl <containing 1-6 C>
G16 = 22-23 13-1 / **bond**

$\text{G11} \text{---} \text{C}$
22 13
 ||
 G9



G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
G18 = O / S / S(O) / SO2 / NH / 33

$\text{N} \text{---} \text{G10}$
33


G19 = carbocycle <containing 5-10 C, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G20) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms), mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G20) /
(Specifically claimed: 150 / 162 / 166)




G20 = alkyl <containing 1-6 C> / F / Cl / Br / I / CN /
 NO2 / SCF3 / CF3 / OCF3 / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH / SH / 35 /
 37 / 40 / 42 / 46 / 47 / 54 / 59 /
 carbocycle <containing 5-6 C, 5- to 6-membered monocyclic
 ring> (opt. substd. by 1 or more G28) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G28) / 61



G21 = O / S / S(O) / SO2
 G22 = alkyl <containing 1-6 C>
 G23 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G24 = (1-3) CH2
 G25 = O / S / S(O) / SO2 / NH / 51



G26 = (0-3) CH2
 G27 = OH / alkoxy <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G28 = alkyl <containing 1-6 C> / F / Cl / Br / I / OH /
 NH2
 G29 = CH2 / O / S / S(O) / SO2 / NH / 63




G30 = carbocycle <containing 5-6 C,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G28) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,

zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring
 (opt. substd. by 1 or more G28)

- G31 = H / alkyl <containing 1 or more C> (opt. substd.) /
 alkenyl <containing 3 or more C> (opt. substd.) /
 alkynyl <containing 3 or more C> (opt. substd.) / 66 /
 carbocycle <containing 5-10 C, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
 (Specifically claimed: 177)

G32-G36
 66 67








- G32 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G33) / 69-9 71-67 /
 (Specifically claimed: CH₂)

G34-G35-G34
 69 71


- G33 = Ph / F / Cl / Br / I / NH₂ / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / CO₂H /
 alkoxy carbonyl <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G35 = O / S / S(O) / SO₂ / NH / 72 / C(O)

N-G10
 72


- G36 = carbocycle <containing 5-10 C, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37)
 G37 = alkyl <containing 1-6 C> / F / Cl / Br / I / CN /
 NO₂ / CF₃ / OCF₃ / NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / 74 / 76 / 81 /
 86 / OH / 94 / SH / 89 / 91 / 96 / 100 / 104 / 105 / 109 /
 112 / 114 / 118 / 136 / 139 / carbocycle <containing 5-6 C,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) / 142 / 144




G38 = NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G39 = NH / 79



G40 = H / alkyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C>
 G41 = 83 / alkylsulfonyl <containing 1-6 C>



G42 = (1-4) CH₂
 G43 = alkyl <containing 1-6 C> / 124 / 126 / 132



G44 = alkylene <containing 1-6 C>
 G45 = cycloalkylene <containing 3-6 C>
 G46 = OH / alkoxy <containing 1-6 C>
 G47 = alkyl <containing 1-6 C> / F / Cl / Br / I / OH /
 CN / tetrazolyl / NH₂ / CO₂H / alkoxycarbonyl <containing
 1-6 C>
 G48 = carbocycle <containing 5-6 C>,

5- to 6-membered monocyclic ring> (opt. substd.) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring> (opt. substd.)
 G49 = carbocycle <containing 5-6 C,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) / 146

G48=O
 146

G50 = CO₂H / H / CO₂Me / CN
 Patent location: claim 1
 Note: substitution is restricted
 Note: additional interruptions in G14 and G32 also
 claimed
 Note: and pharmaceutically acceptable salts and N-oxides
 Stereochemistry: and isomers and racemic forms



W L86 ANSWER 14 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 140:407108 MARPAT
 TITLE: Process for synthesizing antifolates
 INVENTOR(S): Xiao, Zejun; Kochat, Harry
 PATENT ASSIGNEE(S): USA
 SOURCE: U.S. Pat. Appl. Publ., 7 pp.
 CODEN: USXXCO
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004092739	A1	20040513	US 2003-627485	20030725
WO 2004045500	A2	20040603	WO 2003-US33237	20031022
WO 2004045500	A3	20040826		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM,
 PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,
 TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG


AU 2003287174 A1 20040615 AU 2003-287174 20031022
 PRIORITY APPLN. INFO.: US 2002-425826P 20021113
 US 2003-627485 20030725
 WO 2003-US33237 20031022

GI



AB This invention relates to a process for synthesizing certain folic acid analogs [I; herein R1, R2 = amino or N-alkyl substituted amino, HO, alkoxy, keto, lower alkyl, or a nitrogen or oxygen protecting group; R3 = H, HO, alkoxy, CF3, alkoxy, halo, SH, or alkylthio; R4 = HO, alkoxy, CO-X; wherein X = HO, alkoxy, or an amino acid residue; X1-X4 = carbon or nitrogen], in particular γ -methylene glutamate 5,8,10-trideazaaminopterin (TRIDAM), (II) which are useful in treating cancer, inflammatory diseases, or autoimmune diseases, and are commonly referred to as antifolates (no data). The process employs improved steps for annulation, derivatization and addition reactions to produce the described antifolates from commonly available starting materials. Thus, a mixture of 2-amino-5-methylbenzonitrile and cyanoguanidine in 1 N aqueous HCl solution was heated at reflux for 1.5 h to give, after workup and treatment with aqueous ammonium hydroxide, 2,4-diamino-6-methylquinazoline which was amidated with benzoyl chloride in the presence of Et₃N in 1,4-dioxane under heating at reflux for 1 h to give 2,4-dibenzamido-6-methylquinazoline (III). III was brominated by 1,3-dibromo-5,5-dimethylimidazolidine-2,4-dione in the presence of benzoyl peroxide in CCl₄ under irradiation with a high intensity lamp (600 W, 120 V) for 1 h to give 2,4-dibenzamido-6-bromomethylquinazoline which was reacted with triphenylphosphine in THF under relaxing for 2 h and underwent Wittig reaction with Me 4-formylbenzoate in the presence of potassium tert-butoxide in THF at 25° for 24 h to give 2,4-Dibenzamido-6-[2-(p-methoxycarbonylphenyl)ethenyl]quinazoline (IV). IV was hydrogenated over 10% Pd-C in DMF at a hydrogen pressure of 20 psi for 20 h to give 2,4-Dibenzamido-6-[p-(methoxycarbonyl)phenethyl]quinazoline which was hydrolyzed in a mixture of 1 N aqueous KOH solution and MeCN under heating at reflux for 42 h and neutralized with AcOH to give 4-amino-4-deoxy-5,8,10-trideazapteroic acid (V). V was condensed with di-Et 4-methylene-L-glutamate hydrochloride in DMF at 25° for 30 min using 1-hydroxybenzotriazole and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride as condensing agents to give di-Et 4-methylene-N-[4-[2-(2,4-diaminoquinazolin-6-yl)ethyl]benzoyl]glutamate, i.e. TRIDAM di-Et ester, which was saponified in a mixture of 1 N aqueous NaOH solution and MeCN at 25° for 16 h and neutralized with AcOH to give TRIDAM II.

MSTR 1



G1 = NH₂ (opt. substd.) / OH / alkoxy / loweralkyl /
R <"keto group"> / 22 / (Example: NHCOPh)

G2—G3
22



G2 = O / NH
G3 = R <"protecting group">
G4 = N / 24

C—G5
24

G5 = H / OH / alkoxy / CF₃ / alkoxy / halo / SH /
alkylthio
G6 = OH / alkoxy / 26 / (Example: OMe)

C(O)·G7
26

G7 = OH / alkoxy / R <"amino acid residue"> /
(Examples: 29 / 36)



G8 = CH / N
G9 = C / 28

N⁺
28

Patent location: claim 1

L86 ANSWER 15 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 141:350180 MARPAT
TITLE: Preparation of benzoxazinones and benzodioxanes as
antibacterials.

INVENTOR(S): Surivet, Jean-Philippe; Zumbrunn, Cornelia;
Hubschwerlen, Christian
PATENT ASSIGNEE(S): Morphochem A.-G. Aktiengesellschaft fuer
Kombinatorische Chemie, Germany
SOURCE: Ger. Offen., 16 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10316081	A1	20041021	DE 2003-10316081	20030408
AU 2004228147	A1	20041021	AU 2004-228147	20040329
CA 2534891	AA	20041021	CA 2004-2534891	20040329
WO 2004089947	A2	20041021	WO 2004-EP3306	20040329
WO 2004089947	A3	20050106		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG


EP 1613624	A2	20060111	EP 2004-724014	20040329
------------	----	----------	----------------	----------

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK

PRIORITY APPLN. INFO.:


DE 2003-10316081	20030408
WO 2004-EP3306	20040329

GI




AB Title compds. (I; A = O, S, NH, alkylene, alkenylene, alkynylene, heteroalkylene; X1-X5 = N, CH, CR₄; Cy = cycloalkylene, heterocycloalkylene, arylene, heteroarylene; R₁ = H, halo, OH, amino, SH, alkyl, alkoxy, cycloalkyl, etc.; R₂ = halo, OH, amino, NO₂, SH, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, etc.; R₂R₂ = atoms to form aryl, heteroaryl, cycloalkyl, heterocycloalkyl; etc. rings; R₃ = alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, etc.; R₄ = halo, OH, alkyl, alkenyl, alkynyl, heteroalkyl; m, n = 0-2), were prepared as antibacterials (no data). Thus, 4-(7-methoxynaphthalen-1-ylooxymethyl)piperidine (preparation given), 6-(2-chloroacetyl)-4H-benzo[1,4]oxazin-3-one, and Et₃N were heated 2 h at 50° in THF to give 52% coupling product, which was reduced with NaBH₄ in EtOH to give 6-[1-hydroxy-2-[4-(7-methoxynaphthalen-1-ylooxymethyl)piperidin-1-yl]ethyl]-4H-benzo[1,4]dioxane.


MSTR 1



G1 = N / 11






G2 = F / Cl / Br / I / H / OH /
carbon chain <containing 1-20 C, 0 or more double bonds,
0 or more triple bonds> (opt. substd. by G3) /
R <"heteroalkyl group"> / (Specifically claimed: alkoxy
<containing 1-4 C> (opt. substd. by 1 or more F) / 117)




G3 = F / Cl / Br / I

G4 = O / S / NH / carbon chain <containing 1-20 C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by G3) / R <"heteroalkylene group"> /
16-7 17-14 / 18-7 19-14 / 20-7 22-14 /
(Specifically claimed: G6 / 23-7 24-14 / 26-7 25-14 /
CHOH / 29-7 30-14 / 33-7 32-14 / 35-7 36-14 /
37-7 38-14 / 39-7 40-14 / 41-7 42-14 / 43-7 47-14 /
49-7 51-14 / 52-7 58-14 / 54-7 59-14 / 56-7 60-14)






- G5 = O / S / NH
 G6 = (1-2) CH2
 G7 = carbon chain <containing 1-20 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by G3) / R <"heteroalkylene group">
 G8 = O / S / 27



- G9 = alkyl <containing 1-4 C>
 G10 = (1-3) CH2
 G11 = cycloalkylene <containing 3-14 C, non-aromatic>
 (opt. substd. by (up to 2) G12) /
 heterocycle <containing 3-14 atoms, zero or more O,
 zero or more N, zero or more Si, zero or more Se,
 zero or more S, zero or more P (no other heteroatoms),
 non-aromatic> (opt. substd. by (up to 2) G12) /
 arylene <containing 6-14 C> (opt. substd. by (up to 2) G12) /
 heteroarylene <containing 5-14 atoms, zero or more O,
 zero or more N, zero or more P,
 zero or more S (no other heteroatoms)>
 (opt. substd. by (up to 2) G12) /
 (Specifically claimed: 66-13 63-15 / 74-13 70-15)



- G12 = F / Cl / Br / I / OH / NH2 / NO2 / SH /
 carbon chain <containing 1-20 C, 0 or more double bonds,
 0 or more triple bonds> (opt. substd. by G3) /
 R <"heteroalkyl group"> / aryl <containing 6-14 C>
 (opt. substd. by G14) / heteroaryl <containing 5-14 atoms,
 zero or more N, zero or more O, zero or more S,
 zero or more P (no other heteroatoms)>
 (opt. substd. by G14) / carbocycle <containing 3-14 C,
 non-aromatic> (opt. substd. by G13) /
 heterocycle <containing 3-14 atoms, zero or more O,
 zero or more N, zero or more Si, zero or more Se,
 zero or more P, zero or more S (no other heteroatoms),
 non-aromatic> (opt. substd.)
 G13 = R / carbon chain <containing 1-20 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by G3) / R <"heteroalkyl residue">
 G14 = R / carbon chain <containing 1-20 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by G3) / carbocycle <containing 3-14 C,
 non-aromatic> (opt. substd.)
 G15 = N / 67

^C₆₇—G16


G16 = H / OH
 G17 = N / CH
 G18 = (0-1) CH2
 G19 = carbon chain <containing 1-20 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by G3) / R <"heteroalkyl group"> /
 aryl <containing 6-14 C> (opt. substd. by G14) /
 heteroaryl <containing 5-14 atoms, zero or more N,
 zero or more O, zero or more S,
 zero or more P (no other heteroatoms)>
 (opt. substd. by G14) / carbocycle <containing 3-14 C,
 non-aromatic> (opt. substd. by G13) /
 heterocycle <containing 3-14 atoms, zero or more O,
 zero or more N, zero or more Si, zero or more Se,
 zero or more P, zero or more S (no other heteroatoms),
 non-aromatic> (opt. substd.) / (Specifically claimed: 78)

^{G22-G23}₇₈


G20 = H / F / Cl / Br / I / OH / **NH2** / SH /
 alkyl <containing 1-20 C> (opt. substd.) /
 R <"heteroalkyl group"> / 76 / carbocycle <containing 3-14
 C, non-aromatic> (opt. substd. by G13) /
 heterocycle <containing 3-14 atoms, zero or more O,
 zero or more N, zero or more Si, zero or more Se,
 zero or more P, zero or more S (no other heteroatoms),
 non-aromatic> (opt. substd.) / (Specifically claimed: OMe)

^O₇₆—G21


G21 = alkyl <containing 1-20 C> (opt. substd. by G3) /
 R <"heteroalkyl group"> / carbocycle <containing 3-14 C,
 non-aromatic> (opt. substd. by G13) /
 heterocycle <containing 3-14 atoms, zero or more O,
 zero or more N, zero or more Si, zero or more Se,
 zero or more P, zero or more S (no other heteroatoms),
 non-aromatic> (opt. substd.)
 G22 = carbon chain <containing 1-20 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by G3) / R <"heteroalkylene group">
 G23 = aryl <containing 6-14 C> (opt. substd. by G14) /
 heteroaryl <containing 5-14 atoms, zero or more N,
 zero or more O, zero or more S,
 zero or more P (no other heteroatoms)>
 (opt. substd. by G14) / carbocycle <containing 3-14 C,
 non-aromatic> (opt. substd. by G13) /
 heterocycle <containing 3-14 atoms, zero or more N,
 zero or more O, zero or more S, zero or more P,
 zero or more Si, zero or more Se (no other heteroatoms),
 non-aromatic> (opt. substd.) / 80 / 97




G24 = N / 90



G25 = H / F / Cl / Br / I / OH /
carbon chain <containing 1-20 C, 0 or more double bonds,
0 or more triple bonds> (opt. substd. by G3) /
R <"heteroalkyl group">
G26 = (0-2) 93



G27 = O / S / 109



G28 = H / carbon chain <containing 1-20 C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by G3) / R <"heteroalkyl residue">
G29 = alkyl <containing 1-5 C>
Patent location: claim 1

yes
L86 ANSWER 16 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 139:164803 MARPAT
TITLE: Preparation of condensed heterocyclic compounds as
PARP inhibitors
INVENTOR(S): Ishida, Junya; Hattori, Kouji; Kido, Yoshiyuki;
Yamamoto, Hirofumi
PATENT ASSIGNEE(S): Fujisawa Pharmaceutical Co., Ltd., Japan
SOURCE: PCT Int. Appl., 64 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003063874	A1	20030807	WO 2003-JP708	20030127
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL,				

G11 = 27 / (1-3) N

$\overset{\text{C}}{\underset{27}{\text{---}}} \text{G12}$

G12 = H / R / (Specifically claimed: alkoxy <containing
1-4 C> / F / Cl / Br / I / NH₂ /
alkylamino <containing 1-4 C> /
dialkylamino <each alkyl containing 1-4 C> / NHNH₂ /
alkyl <containing 1-4 C> / OH)

G13 = 38 / (up to 1) N

$\overset{\text{C}}{\underset{38}{\text{---}}} \text{G12}$

G14 = NH / O / S
G15 = 58 / (up to 1) N

$\overset{\text{C}}{\underset{58}{\text{---}}} \text{G12}$

G16 = 60 / (up to 1) N

$\overset{\text{C}}{\underset{60}{\text{---}}} \text{G12}$

G17 = (1-2) CH₂
G18 = N / CH
G19 = O / S / NH / 149

$\overset{\text{N}}{\underset{149}{\text{---}}} \text{G2}$


G20 = H / alkyl <containing 1-4 C> / NH₂ /
alkylamino <containing 1-4 C> /
dialkylamino <each alkyl containing 1-4 C>

Derivative: or pharmaceutical salts or hydrates
Patent location: claim 1
Note: substitution is restricted; additional ring
formation is allowed; also incorporates claim 75
Stereochemistry: and stereoisomers


L86 ANSWER 26 OF 26 BEILSTEIN COPYRIGHT 2006 BEILSTEIN MDL on STN

Beilstein Records (BRN):	573824
Chemical Name (CN):	3-azido-2-methyl-5-phenyl-2H-pyrimido<4,5-c>quinolin-1-one
Autonom Name (AUN):	3-azido-2-methyl-5-phenyl-2H-pyrimido<4,5-c>quinolin-1-one
Molec. Formula (MF):	C18 H12 N6 O











alkoxycarbonyl <containing 1-4 C> /
 cycloalkyloxycarbonyl <containing 3-8 C> /
 alkenyl <containing 2-4 C> / cycloalkenyl <containing 4-8 C>
 / alkynyl <containing 2-4 C>



G₇ = cycloalkyl <containing 3-8 C>
 G₈ = SO₂ / C(O)
 G₉ = NH₂ / alkylamino <containing 1-4 C> /
 dialkylamino <each alkyl containing 1-4 C> /
 cycloalkylamino <containing 3-8 C> / 14



G₁₀ = 22 / 34 / 45 / 54 / (Specifically claimed: 71 / 80 /
 94 / 114 / 123 / 138)



(dimethylamino)pyrido[3,4-d]pyrimidine, which demonstrated a IC50 of 6 pM for inhibition of tyrosine kinase at an epidermal growth factor receptor.

MSTR 1

G10-G1-G5
1 3

G1 = O / S / NH / 5 / 148-1 4-3

G19-G3 N-G2
148 4 5

G2 = alkyl <containing 1-4 C> / OH / NH2 /
alkoxy <containing 1-4 C> / alkylamino <containing 1-4 C>
G3 = (1-2) 7


HC-G4
7

G4 = H / alkyl <containing 1-4 C>
G5 = Ph (opt. substd. by (1-3) G6) / thieryl / furyl /
pyrrolyl / pyridyl / pyrimidinyl / imidazolyl / pyrazinyl /
oxazolyl / thiazolyl / naphthyl / benzothienyl /
benzofuranyl / indolyl / quinoliny / isoquinoliny /
quinazolinyl / carbocycle <containing 10 C, aromatic,
bonds all normalized, bicyclic, (2) 6-membered rings>
(opt. substd. by (1-3) G6) / heterocycle <containing 1-2
heteroatoms, zero or more N, up to 1 O,
up to 1 S (no other heteroatoms), aromatic, 2 double bonds,
5-membered monocyclic ring> (opt. substd. by (1-3) G6) /
heterocycle <containing 1-2 heteroatoms,
1-2 N (no other heteroatoms), aromatic,
bonds all normalized, mono- or bicyclic,
(1-2) 6-membered rings only> (opt. substd. by (1-3) G6) /
heterocycle <containing 1 heteroatom, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
aromatic, 6 normalized bonds, 1 double bond, bicyclic,
(1) 5-membered ring, (1) 6-membered ring>
(opt. substd. by (1-3) G6)
G6 = alkyl <containing 1-4 C> /
cycloalkyl <containing 3-8 C> / alkoxy <containing 1-4 C> /
cycloalkyloxy <containing 3-8 C> / NO2 / F / Cl / Br / I /
perfluoroalkyl <containing 1-4 C> / OH /
alkylcarbonyloxy <containing 1-4 C> /
cycloalkylcarbonyloxy <containing 3-8 C> / NH2 /
alkylamino <containing 1-4 C> /
dialkylamino <each alkyl containing 1-4 C> /
cycloalkylamino <containing 3-8 C> / 9 / CH2OH /
alkylcarbonyl <containing 1-4 C> /
cycloalkylcarbonyl <containing 3-8 C> / CN /
alkylthio <containing 1-4 C> / alkylsulfinyl <containing 1-4
C> / alkylsulfonyl <containing 1-4 C> /
cycloalkylthio <containing 3-8 C> /
cycloalkylsulfinyl <containing 3-8 C> /
cycloalkylsulfonyl <containing 3-8 C> / 12 / SH / CO2H /

PATENT INFORMATION:






PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9519774	A1	19950727	WO 1995-US941	19950123
W: AM, AU, BG, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SI, SK, TJ, UA, UZ				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5654307	A	19970805	US 1994-358351	19941223
ZA 9500440	A	19951010	ZA 1995-440	19950119
AU 9517314	A1	19950808	AU 1995-17314	19950123
AU 686334	B2	19980205		
EP 742717	A1	19961120	EP 1995-909316	19950123
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 09508127	T2	19970819	JP 1995-519732	19950123
PL 179132	B1	20000731	PL 1995-315633	19950123
MD 1632	F2	20010331	MD 1996-217	19950123
RU 2174980	C2	20011020	RU 1996-116985	19950123
RO 117257	B1	20011228	RO 1996-1517	19950123
NZ 281011	A	20020201	NZ 1995-281011	19950123
BG 63245	B1	20010731	BG 1996-100614	19960520
FI 9602856	A	19960925	FI 1996-2856	19960715
FI 114213	B1	20040915		
NO 9603094	A	19960724	NO 1996-3094	19960724
NO 309892	B1	20010417		
FI 2004000648	A	20040507	FI 2004-648	20040507
FI 2004000649	A	20040507	FI 2004-649	20040507
PRIORITY APPLN. INFO.:				US 1994-186735
				US 1994-186745
				US 1994-358351
				WO 1995-US941

GI







I






AB The title compds. [I; A-E = nitrogen with the remaining atom(s) carbon, or any two contiguous positions in A-E taken together can be a single heteroatom N, O or S, in which case one of the two remaining atoms must be carbon, and the other can be either carbon or nitrogen, etc.; A1 = divalent Ph, thienyl, furanyl pyrimidinyl, heterocyclyl, etc.; R1 = H, lower alkyl; R2 = lower alkyl, cycloalkyl, alkoxy, cycloalkoxy, NO2, halogen, etc.; R3-R6 = H, alkyl, alkoxy, HO, acyloxy, (un)substituted NH2, etc.; X = O, S, (un)substituted NH; m = 0-3; n = 0-2], useful for inhibiting tyrosine kinases of the epidermal growth factor receptor family, are prepared Thus, 4-(3-bromoanilino)-6-fluoropyrido[3,4-d]pyrimidine was reacted with Me2NH, producing 4-(3-bromoanilino)-6-



G18 = morpholino / NMe2 / piperazino / pyrrolidino
 G20 = Cl / F / Br / Me / Et / Pr-i / OMe / OH / NH2 /
 CF3 / OCF3 / SMe / OMe / CO2H / CO2Me / CH2NH2 / NMe2 / 116 /
 CH2OH / (Specifically claimed: 124 / NHC(NH)NH2 / 170 / 125)




G21 = morpholino / piperazino / pyrrolidino
 G22 = H / Cl / F / Br / Me / OMe / OH / CF3 / OCF3 /
 OPr-n / NH2 / NMe2 / NHSO2Ph / 149 / 152 / 156 / 158 / 164




Derivative: or pharmaceutically acceptable salts
 Patent location: claim 1
 Note: additional ring formation also claimed


L86 ANSWER 25 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 124:8839 MARPAT
 TITLE: Preparation of bicyclic pyrimidines capable of
 inhibiting tyrosine kinases of the epidermal growth
 factor receptor family
 INVENTOR(S): Bridges, Alexander James; Denny, William Alexander;
 Fry, David; Kraker, Alan; Meyer, Robert; Rewcastle,
 Gordon William; Thompson, Andrew Mark
 PATENT ASSIGNEE(S): Warner-Lambert Co., USA
 SOURCE: PCT Int. Appl., 218 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 4




G9 = NH / 20





G10 = alkyl <containing 1-3 C> (opt. substd.) /
 alkenyl <containing 2-3 C> (opt. substd.) /
 G11 = H / alkyl <containing 1-3 C> (opt. substd.) /
 alkenyl <containing 2-3 C> (opt. substd.) /
 G12 = OH / 46




G13 = N / 48




G14 = H / alkyl <containing 1-3 C> (opt. substd.) /
 alkenyl <containing 2-3 C> (opt. substd.) / NH2 / 50 / 53 /
 OH / 55 / SH / 57 / 59 / CO2H / 66 /
 (Specifically claimed: Me)




G15 = NH2 / 61 / 64



G16 = C(O) / SO2
 G17 = Cl / F / Br / Me / OMe / OH / CF3 / OCF3 / OPr-n /
 NH2 / NMe2 / NHSO2Ph / 87 / 97 / 135 / 99 / 105



G1 = Ph (opt. substd.) / heterocycle <containing 5-6 atoms, aromatic, 5- to 6-membered monocyclic ring> (opt. substd.) / aryl <containing 8-10 C, bicyclic> (opt. substd.) / heterocycle <containing 8-10 atoms, aromatic, bicyclic> (opt. substd.) / (Specifically claimed: Ph (opt. substd. by (1-3) G17) / pyridyl (opt. substd. by (1-3) G17) / 75)




G2 = Ph (opt. substd.) / heterocycle <containing 5-6 atoms, aromatic, 5- to 6-membered monocyclic ring> (opt. substd.) / aryl <containing 8-10 C, bicyclic> (opt. substd.) / heterocycle <containing 8-10 atoms, aromatic, bicyclic> (opt. substd.) / (Specifically claimed: Ph (opt. substd. by (1-3) G20) / pyridyl (opt. substd. by (1-3) G20))

G3 = O / NH (opt. substd.)

G4 = H / alkyl <containing 1-3 C> / OH / alkoxy <containing 1-3 C>


G5 = N / 16



G6 = H / alkyl <containing 1-3 C> / alkenyl <containing 2-3 C> / alkynyl <containing 2-3 C> / Ph (opt. substd. by (1-3) G7)

G7 = halo / OMe / CN / NO2 / NH2 / OH / Me / Et

G8 = S / O / SO2 / S(O) / 18-9 19-14 / 23-9 22-14 / 25-9 26-14 / 29-9 28-14 / C(O) / 30-9 31-14 / 33-9 32-14 / 34-9 35-14 / 37-9 36-14 / NH / 38 / 41 / 44



TITLE: Heterocyclic compound inhibitors of p38 kinase, pharmaceutical compositions, and therapeutic use
 INVENTOR(S): Salituro, Francesco; Bemis, Guy; Cochran, John
 PATENT ASSIGNEE(S): Vertex Pharmaceuticals Incorporated, USA
 SOURCE: PCT Int. Appl., 68 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9964400	A1	19991216	WO 1999-US12951	19990611
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9944297	A1	19991230	AU 1999-44297	19990611
EP 1086085	A1	20010328	EP 1999-927377	19990611
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
EP 1277740	A1	20030122	EP 2002-22891	19990611
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
US 2001025044	A1	20010927	US 2000-734069	20001211
US 6528508	B2	20030304		
US 2003149037	A1	20030807	US 2002-327020	20021220
US 6800626	B2	20041005		
US 2005049251	A1	20050303	US 2004-951409	20040927

PRIORITY APPLN. INFO.:

US 1998-89147P 19980612
 EP 1999-927377 19990611
 WO 1999-US12951 19990611
 US 2000-734069 20001211
 US 2002-327020 20021220

AB The invention relates to heterocyclic compound inhibitors of p38, a mammalian protein kinase involved cell proliferation, cell death and response to extracellular stimuli. The invention also relates to methods for producing these inhibitors. The invention also provides pharmaceutical compns. comprising the inhibitors of the invention and methods of utilizing those compns. in the treatment and prevention of various disorders.










REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1A





G15 = 88 / 89



G16 = heteroaryl (opt. substd. by (1-3) G17) /
 aryl (opt. substd. by (1-3) G17) /
 (Specifically claimed: 159) / (Examples: Ph / naphthyl /
 pyridyl / thienyl / furyl / pyrimidinyl / pyrazinyl / 172 /
 180 / 189 / 200 / 210 / 217 / 222 / 231)




G17 = F / Cl / Br / alkyl <containing 1-5 C>
 (opt. substd. by OH) / alkoxy <containing 1-4 C> /
 alkyl <containing 1-5 C> (substd. by (1-3) G10) / OH / SH /
 alkylthio <containing 1-4 C> / alkylcarbonyl <containing 1-4
 C> / NH₂ / 147 / 149 / alkyl <containing 1-4 C>
 (substd. by alkoxy <containing 1-4 C>) /
 alkyl <containing 1-4 C> (substd. by NH₂) /
 alkyl <containing 1-5 C> (substd. by 152) /
 alkyl <containing 1-5 C> (substd. by 154) / NO₂




Patent location: claim 1


L86 ANSWER 24 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 132:30857 MARPAT




G9 = H / F / Cl / Br / alkyl <containing 1-5 C> (opt. substd. by OH) / cycloalkyl <containing 3-8 C> / alkoxy <containing 1-4 C> / alkylcarbonyl <containing 1-4 C> / aryl (opt. substd.) / heteroaryl / alkylthio <containing 1-4 C> / alkyl <containing 1-5 C> (substd. by (1-3) G10) / alkyl <containing 1-4 C> (substd. by alkoxy <containing 1-4 C>) / alkyl <containing 1-5 C> (substd. by 43) / alkyl <containing 1-5 C> (substd. by 45) / NO₂ / (Specifically claimed: Me)




G10 = F / Cl / Br
 G11 = CH=CHCH=CH / CH₂CH₂CH₂CH₂
 G12 = H / alkyl <containing 1-10 C> / alkyl <containing 1-4 C> (substd. by cycloalkyl <containing 3-8 C>) / cycloalkyl <containing 3-8 C>
 G13 = 76 / N




G14 = C(O) / 85











G6 = alkyl <containing 1-10 C> /
 alkyl <containing 1-4 C> (substd. by cycloalkyl <containing 3-8 C>) / cycloalkyl <containing 3-8 C> /
 aryl (opt. substd.) / heteroaryl /
 alkyl <containing 1-5 C> (substd. by OH) /
 alkyl <containing 1-5 C> (substd. by 1 or more aryl (opt. substd.)) / alkyl <containing 1-5 C>
 (substd. by heteroaryl) / alkylcarbonyl <containing 1-4 C> /
 alkyl <containing 1-4 C> (substd. by alkoxy <containing 1-4 C>) / alkyl <containing 1-4 C> (substd. by alkylthio
 <containing 1-4 C>) / alkyl <containing 1-4 C>
 (substd. by NH2) / alkyl <containing 1-5 C> (substd. by 20) /
 alkyl <containing 1-5 C> (substd. by G7) /
 alkyl <containing 1-3 C> (substd. by CO2H) /
 alkyl <containing 1-3 C> (substd. by alkoxycarbonyl
 <containing 1-3 C>) / (Specifically claimed: Et / Bu-n /
 Pr-n / 169)



G7 = 22 / heterocycle <containing 1 heteroatom, 1 N,
 attached through 1 N, 4- to 6-membered monocyclic ring>



G8 = 30-4 26-2 27-25 / 34-4 35-2 31-25 /
 36-4 37-2 38-25 / 49-4 50-2 51-25 / 56-4 57-2 58-25 /
 65-4 66-2 62-25 / 72-4 73-2 74-25 / 81-4 82-2 83-25 /
 94-4 95-2 96-25 / 103-4 104-2 105-25 /
 110-4 111-2 112-25 / 118-4 119-2 120-25 /
 126-4 127-2 128-25 / 130-4 131-2 132-25 /
 139-4 135-2 136-25 / 143-4 144-2 140-25




PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000038350	A2	20000208	JP 1999-136173	19990517
PRIORITY APPLN. INFO.:			JP 1998-135673	19980518

AB Pyrimidine derivs. (Markush's structures given), including 4-(N-butyl-N-ethylamino)-2,5-di-methyl-7-(2,4,6-trimethylphenyl)-thieno[3,4-d]pyrimidine, and their pharmaceutically acceptable salts and hydrates are claimed as antidiabetics by acting as CRF antagonists. The hypoglycemic, insulin secretion-promoting, and insulin-enhancing effects were tested, and a formulation example of tablets was given.

MSTR 1





G1 = N / 6




G2 = H / alkyl <containing 1-5 C> / F / Cl / Br / CN / OH / alkoxy <containing 1-4 C>

G3 = H / alkyl <containing 1-5 C> (opt. substd. by OH) / alkyl <containing 1-4 C> (substd. by cycloalkyl <containing 3-8 C>) / cycloalkyl <containing 3-8 C> / alkoxy <containing 1-4 C> / NH2 / 9 / 11 / alkyl <containing 1-5 C> (substd. by (1-3) G10) / alkyl <containing 1-5 C> (substd. by 1 or more aryl (opt. substd.)) / alkyl <containing 1-4 C> (substd. by alkoxy <containing 1-4 C>) / OH / aryl (opt. substd.) / heteroaryl / F / Cl / Br / SH / alkylthio <containing 1-4 C> / (Specifically claimed: Me)




G4 = alkyl <containing 1-5 C> / cycloalkyl <containing 3-8 C>


G5 = NH2 / 15 / 17 / heterocycle <containing 1 or more heteroatoms, 1 or more N, attached through 1 or more N> / alkoxy <containing 1-10 C> / alkoxy <containing 1-4 C> (substd. by cycloalkyl <containing 3-8 C>) / cycloalkyloxy <containing 3-8 C>

G23-G10
127

G18 = alkylene <containing 1 or more C> (opt. substd.)
 G19 = O / NH
 G20 = 2-pyridyl / Ph / 176



G21 = Ph (opt. substd.) / naphthyl / biphenyl / 183 /
 189 / 201 / carbocycle <containing 5-10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G12) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more O, zero or more S,
 zero or more N (no other heteroatoms), 1 or more C,
 mono- or bicyclic> / heterocycle <containing 5-10 atoms,
 1-4 heteroatoms, 1-4 N (no other heteroatoms), 1 or more C,
 mono- or bicyclic> (substd. by 1 or more G12) /
 alkyl <containing 1 or more C> (opt. substd.) / CF₃ /
 cycloalkyl <containing 3-7 C> (opt. substd. by aryl
 <containing 6-14 C>) / cycloalkyl <containing 7-16 C,
 2-3 rings>



G22 = R / F
 G23 = R <"linking group"> / alkylene <containing 1 or
 more C> (opt. substd.)
 Patent location: claim 1
 Note: or pharmaceutically acceptable salts
 Note: substitution is restricted

L86 ANSWER 23 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 132:146643 MARPAT
 TITLE: Pyrimidine CRF antagonists as antidiabetics
 INVENTOR(S): Seio, Yasushi; Tanaka, Hiroshi; Goto, Shinji; Amano,
 Yusaku
 PATENT ASSIGNEE(S): Yoshitomi Pharmaceutical Industries, Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 24 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1




G14 = zero or more N (no other heteroatoms) >
= NH2 / 99

HN—G15
99

G15 = Ph (opt. substd. by 1 or more G11) / naphthyl / biphenyl / 107 / 113 / 125 / carbocycle <containing 5-10 C, mono- or bicyclic> (opt. substd. by 1 or more G12) / heterocycle <containing 5-10 atoms, 1-4 heteroatoms, zero or more O, zero or more S, zero or more N (no other heteroatoms), 1 or more C, mono- or bicyclic> / heterocycle <containing 5-10 atoms, 1-4 heteroatoms, 1-4 N (no other heteroatoms), 1 or more C, mono- or bicyclic> (substd. by 1 or more G12) / alkyl <containing 1-7 C> (opt. substd. by 1 or more G13) / CF3 / cycloalkyl <containing 3-7 C> (opt. substd. by aryl <containing 6-14 C>) / cycloalkyl <containing 7-16 C, 2-3 rings> / 101 / 103 / 150 / 152 / 154 / (Specifically claimed: 165 / 173 / CH2CH2Ph)

G16-G10
101



C(O)-G19-G17
103




C(O)-G17
150

O2S—G17
152




C(O)NH—G18-G17
154



G16 = R <"linking group"> / alkylene <containing 1 or more C> (opt. substd. by 1 or more G22)
G17 = Ph (opt. substd.) / naphthyl / biphenyl / 130 / 136 / 148 / carbocycle <containing 5-10 C, mono- or bicyclic> (opt. substd. by 1 or more G12) / heterocycle <containing 5-10 atoms, 1-4 heteroatoms, zero or more O, zero or more S, zero or more N (no other heteroatoms), 1 or more C, mono- or bicyclic> / heterocycle <containing 5-10 atoms, 1-4 heteroatoms, 1-4 N (no other heteroatoms), 1 or more C, mono- or bicyclic> (substd. by 1 or more G12) / alkyl <containing 1 or more C> (opt. substd.) / CF3 / cycloalkyl <containing 3-7 C> (opt. substd. by aryl <containing 6-14 C>) / cycloalkyl <containing 7-16 C, 2-3 rings> / 127

G14-G6
32 33

G6 = 34-32 38-2 / 40-32 44-2 / 46-32 50-2






G7 = H / F / Cl / Br / I / alkyl <containing 1-4 C> /
cycloalkyl <containing 3-7 C> / CF3 /
(Specifically claimed: Me / Pr-i / Et)

G8 = H / F / Cl / Br / I

G9 = CH2 / S / SO2

G10 = Ph (opt. substd. by 1 or more G11) / naphthyl /
biphenyl / 79 / 85 / 97 / carbocycle <containing 5-10 C,
mono- or bicyclic> (opt. substd. by 1 or more G12) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more O, zero or more S,
zero or more N (no other heteroatoms), 1 or more C,
mono- or bicyclic> / heterocycle <containing 5-10 atoms,
1-4 heteroatoms, 1-4 N (no other heteroatoms), 1 or more C,
mono- or bicyclic> (substd. by 1 or more G12) /
alkyl <containing 1-7 C> (opt. substd. by 1 or more G13) /
CF3 / cycloalkyl <containing 3-7 C>
(opt. substd. by aryl <containing 6-14 C>) /
cycloalkyl <containing 7-16 C, 2-3 rings>



G11 = alkyl <containing 1-4 C> /
alkoxy <containing 1-4 C> / F / Cl / Br / I / OH / CO2H /
CONH2 / CH2OH / alkoxy carbonyl <containing 1-4 C> / SO2NH2

G12 = F / Cl / Br / I / OH


G13 = OH / CO2H / NH2 / aryl <containing 6-14 C> /
cycloalkyl <containing 3-7 C> / CF3 / NMe2 /
alkyl <containing 1-3 C> (substd. by 1 or more aryl
<containing 6-14 C>) / heteroaryl <containing 5-10 atoms,
zero or more O, zero or more S,
zero or more N (no other heteroatoms)> /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more O, zero or more S,

SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
 YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 CA 2403558 AA 20010927 CA 2001-2403558 20010319
 EP 1267876 A1 20030102 EP 2001-918822 20010319
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
 JP 2003527426 T2 20030916 JP 2001-568427 20010319
 US 2002006923 A1 20020117 US 2001-815404 20010322
 US 6534510 B2 20030318
 PRIORITY APPLN. INFO.: US 2000-191559P 20000323
 WO 2001-US8733 20010319
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Heterocyclic compds. I (T = N or CY1, NH or CY1Y2 and N = 0, 1, 2 and Y1 and Y2 are independently H, C1-4 alkyl, halogen, NH2, OH; A = a-d; W = H, R1, R1OCO, R1CO, R1SO2, R1(CH2)nNHCO, (R1)2CH(CH2)nNHCO where n = 0-4 and R1 = (non)heterocyclic; R4 = Ph (un)substituted with one or more C1-4 alkyl(alkoxy), halogen, OH, CO2H, CONH2, CH2OH, naphthyl, biphenyl, pyridine N-oxide, (un)saturated (non)heterobicyclic ring, C1-7 alkyl (un)substituted with OH, CO2H, NH2, CF3, NMe2, etc.; R3, R5, and R6 = independently H, halogen, C1-4 alkyl, C3-7 cycloalkyl, CF3; X = H, halogen; Z = CH2, S, SO2) were prepared as thrombin inhibitors (no data). Thus II was prepared from 8-aminomethyl-1,2,3,4-tetrahydronaphthyridine and III in DMF, HOAt and EDC. Oral dosages of the thrombin inhibitors will range between 0.01 mg/kg of body weight per day to about 30 mg/kg/day.
 REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1A



G2 = N / 27

$\overset{C}{\underset{27}{\text{---}}}G3$

G3 = H / alkyl <containing 1-4 C> / F / Cl / Br / I /
 NH2 / OH
 G5 = 33 / 63

G16 = 19 / N

$$\begin{array}{c} \text{C} \text{---} \text{G4} \\ 19 \end{array}$$

G17 = S / S(O) / SO2

G18 = NH2 / 36 / heterocycle <containing 1 or more N, zero or more O, zero or more S (no other heteroatoms), attached through 1 N, mono- or polycyclic>

$$\begin{array}{c} \text{G9} \text{---} \text{G8} \\ 36 \end{array}$$

G19 = OH / 40 / NH2 / 42 / heterocycle <containing 1 or more N, zero or more O, zero or more S (no other heteroatoms), attached through 1 N, mono- or polycyclic>

$$\begin{array}{cc} \text{O} \text{---} \text{G7} & \text{G9} \text{---} \text{G8} \\ 40 & 42 \end{array}$$

G20 = OH / 46

$$\begin{array}{c} \text{O} \text{---} \text{G7} \\ 46 \end{array}$$

G21 = O / S / S(O) / SO2

G22 = SO2 / C(O)

Patent location:

claim 1

Note:

and tautomers, and pharmaceutically acceptable salts, prodrugs and solvates

Note:

additional ring formation also claimed

Note:

additional oxo formation also claimed

Stereochemistry:

and isomers, enantiomers, diastereomers

L86 ANSWER 22 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 135:272974 MARPAT

TITLE: Preparation of heterocyclic compounds as thrombin inhibitors

INVENTOR(S): Barrow, James C.; Dorsey, Bruce D.; Selnick, Harold G.; Ngo, Phung L.

PATENT ASSIGNEE(S): Merck & Co., Inc., USA

SOURCE: PCT Int. Appl., 84 pp.

CODEN: PIXXD2

DOCUMENT TYPE:

Patent

LANGUAGE:

English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001070229	A1	20010927	WO 2001-US8733	20010319
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,				

G3 = H / OH / halo / CN / 44 / NH2 / 48 /
 heterocycle <containing 1 or more N, zero or more O,
 zero or more S (no other heteroatoms), attached through 1 N,
 mono- or polycyclic> / alkyl <containing 1-12 C>
 (opt. substd.) / alkenyl <containing 2-12 C> (opt. substd.) /
 alkynyl <containing 2-12 C> (opt. substd.) /
 carbocycle <containing 3 or more C, 0 or more double bonds,
 mono- or polycyclic> (opt. substd.) /
 heterocycle <containing 3 or more atoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or polycyclic>

$\overset{\text{C(O)-G20}}{\underset{44}{\text{C}}}$ $\overset{\text{G9-G8}}{\underset{48}{\text{S}}}$

G4 = H / **halo** / NO2 / CN / 50 / OH / SH / 53 / 55 / NH2 /
 57 / heterocycle <containing 1 or more N, zero or more O,
 zero or more S (no other heteroatoms), attached through 1 N,
 mono- or polycyclic> / 59 / alkylcarbonyl <containing 1-12 C>
 (opt. substd.) / carbon chain <containing 1-12 C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)

$\overset{\text{G21-G7}}{\underset{50}{\text{O}}}$ $\overset{\text{O}}{\underset{\text{S}-\text{OH}}{\text{S}}}$ $\overset{\text{S}-\text{OH}}{\underset{55}{\text{S}}}$ $\overset{\text{G9-G8}}{\underset{57}{\text{S}}}$ $\overset{\text{G22-G19}}{\underset{59}{\text{S}}}$

G7 = R
 G8 = R
 G9 = NH / 23

$\overset{\text{N-G8}}{\underset{23}{\text{N}}}$

G10 = R
 G11 = **C(O)** / S(O) / SO2
 G12 = 13 / **N**



$\overset{\text{C-G3}}{\underset{13}{\text{C}}}$

G13 = **NH** / O / S
 G14 = 15 / N

$\overset{\text{C-G4}}{\underset{15}{\text{C}}}$


G15 = 17 / **N**

$\overset{\text{C-G4}}{\underset{17}{\text{C}}}$



AB Title compds. I [wherein X1 = CO, SO, or SO₂; X2 = CR₃ or N; X3 = NH, O, or S; X4 = CR₄ or N; X5 = CR₅ or N; X6 = CR₆ or N] were prepared were prepared as inosine monophosphate dehydrogenase (IMPDH) enzyme inhibitors. For example, acetalization of 4-nitro-2-methoxytoluene with AcOH (51%), reduction to the aldehyde (91%), and cycloaddn. with (p-tolylsulfonyl)methyl isocyanate gave 5-(4-nitro-2-methoxyphenyl)oxazole (84%), which was reduced to the amine (95%). Alkylation with Et benzoylacetate and cyclization afforded the 6-(5-oxazolyl)-4(1H)-quinolinone II. Thus, I are useful as therapeutic agents for IMPDH-associated disorders, such as allograft rejection (no data).


MSTR 1



G1 = alkyl <containing 1-12 C> (opt. substd.) /
 alkenyl <containing 2-12 C> (opt. substd.) /
 alkynyl <containing 2-12 C> (opt. substd.) / **NH2** / 21 /
 heterocycle <containing 1 or more N, zero or more O,
 zero or more S (no other heteroatoms), attached through 1 N,
 mono- or polycyclic> / 25 / carbocycle <containing 3 or more
 C, non-aromatic, 0 or more double bonds, mono- or polycyclic>
 (opt. substd.) / aryl <containing 6 or more C,
 mono- or polycyclic> (opt. substd.) /
 heterocycle <containing 3 or more atoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds, mono- or polycyclic> /
 heteroaryl <containing zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or polycyclic>



G2 = F / Cl / Br / I / CN / NO₂ / OH / SH / 32 / 30 /
 27 / 34 / 38 / heteroaryl <containing zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 mono- or polycyclic>



G49 = carbocycle <containing 5-6 C,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G47) / 146















G48=O
 146

Patent location: claim 1
 Note: substitution is restricted
 Note: additional interruptions in G14 and G32 also
 claimed
 Note: and pharmaceutically acceptable salts and N-oxides
 Stereochemistry: and isomers and racemic forms


L86 ANSWER 21 OF 26 MARPAT COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 135:344472 MARPAT
 TITLE: Preparation of 6-(5-oxazolyl)-4(1H)-quinolinones as
 inhibitors of IMPDH enzyme
 INVENTOR(S): Iwanowicz, Edwin J.; Watterson, Scott H.; Dhar, T. G.
 Murali; Pitts, William J.; Gu, Henry H.
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA
 SOURCE: PCT Int. Appl., 263 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001081340	A2	20011101	WO 2001-US12900	20010419
WO 2001081340	A3	20020523		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2407370	AA	20011101	CA 2001-2407370	20010419
EP 1276739	A2	20030122	EP 2001-928708	20010419
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2003531205	T2	20031021	JP 2001-578430	20010419
US 2002040022	A1	20020404	US 2001-840503	20010423
US 6919335	B2	20050719		
PRIORITY APPLN. INFO.:			US 2000-199420P	20000424
			WO 2001-US12900	20010419


GI






G38 = NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C>
G39 = NH / 79



G40 = H / alkyl <containing 1-6 C> / OH /
alkoxy <containing 1-6 C>
G41 = 83 / alkylsulfonyl <containing 1-6 C>



G42 = (1-4) CH₂.
G43 = alkyl <containing 1-6 C> / 124 / 126 / 132



G44 = alkylene <containing 1-6 C>
G45 = cycloalkylene <containing 3-6 C>
G46 = OH / alkoxy <containing 1-6 C>
G47 = alkyl <containing 1-6 C> / F / Cl / Br / I / OH /
CN / tetrazolyl / NH₂ / CO₂H / alkoxycarbonyl <containing
1-6 C>
G48 = carbocycle <containing 5-6 C,
5- to 6-membered monocyclic ring> (opt. substd.) /
heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
5- to 6-membered monocyclic ring> (opt. substd.)

~~G32-G36~~
66 67





G32 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by 1 or more G33) / 69-9 71-67 /
(Specifically claimed: CH2)

~~G34-G35-G34~~
69 71

G33 = Ph / F / Cl / Br / I / NH2 / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C> / CO2H /
alkoxycarbonyl <containing 1-6 C>
G34 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
G35 = O / S / S(O) / SO2 / NH / 72 / C(O)

~~N-G10~~
72

G36 = carbocycle <containing 5-10 C, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms), mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
(Specifically claimed: Ph (opt. substd.) / pyridyl / 186 /
194 / 204 / 212)



G37 = alkyl <containing 1-6 C> / F / Cl / Br / I / CN /
NO2 / CF3 / OCF3 / NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / 74 / 76 / 81 /
86 / OH / 94 / SH / 89 / 91 / 96 / 100 / 104 / 105 / 109 /
112 / 114 / 118 / 136 / 139 / carbocycle <containing 5-6 C,
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G47) /
heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G47) / 142 / 144

~~G24-G38~~
74



~~G39-C(O)-G40~~
76

~~G39-G41~~
81


G41
N-G41
86

S-OH
89


37 / 40 / 42 / 46 / 47 / 54 / 59 /
 carbocycle <containing 5-6 C, 5- to 6-membered monocyclic
 ring> (opt. substd. by 1 or more G28) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G28) / 61



G21 = O / S / S(O) / SO2
 G22 = alkyl <containing 1-6 C>
 G23 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G24 = (1-3) CH2
 G25 = O / S / S(O) / SO2 / NH / 51




G26 = (0-3) CH2
 G27 = OH / alkoxy <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G28 = alkyl <containing 1-6 C> / F / Cl / Br / I / OH /
 NH2
 G29 = CH2 / O / S / S(O) / SO2 / NH / 63




G30 = carbocycle <containing 5-6 C,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G28) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G28)
 G31 = H / alkyl <containing 1 or more C> (opt. substd.) /
 alkenyl <containing 3 or more C> (opt. substd.) /
 alkynyl <containing 3 or more C> (opt. substd.) / 66 /
 carbocycle <containing 5-10 C, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G37)




- G12 = alkyl <containing 1-6 C>
(opt. substd. by 1 or more aryl) / cycloalkyl / aryl /
heteroaryl <containing zero or more N, zero or more O,
zero or more S>
- G13 = H / alkyl <containing 1-6 C>
(opt. substd. by 1 or more aryl) / aryl /
heterocycle <containing zero or more N, zero or more O,
zero or more S> / cycloalkyl
- G14 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by 1 or more G15) / 30-28 32-29







- G15 = F / Cl / Br / I / NH₂ / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C> / CO₂H /
alkoxycarbonyl <containing 1-6 C>
- G16 = 22-23 13-1 / **bond**




- G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
- G18 = O / S / S(O) / SO₂ / NH / 33




- G19 = carbocycle <containing 5-10 C, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G20) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms), mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G20) /
(Specifically claimed: Ph (opt. substd.) / pyridyl / 150 /
158 / 168 / 176)




- G20 = alkyl <containing 1-6 C> / F / Cl / Br / I / CN /
NO₂ / SCF₃ / CF₃ / OCF₃ / NH₂ /
alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH / SH / 35 /




- G1 = H / NH₂ / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G2) /
 alkenyl <containing 3-6 C> (opt. substd. by 1 or more G3) /
 alkynyl <containing 3-6 C> (opt. substd. by 1 or more G3) /
 aryl (opt. substd. by 1 or more G3) /
 heterocycle <containing zero or more N, zero or more O,
 zero or more S> (opt. substd. by 1 or more G3) /
 (Specifically claimed: Me)
- G2 = alkylamino <containing 1-6 C>
 (opt. substd. by 1 or more G3) /
 dialkylamino <each alkyl containing 1-6 C>
 (opt. substd. by 1 or more G3) /
 aryl (opt. substd. by 1 or more G3) /
 cycloalkyl <containing 3-6 C> (opt. substd. by 1 or more G3)
 / NH₂ / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G4) / CN / CO₂H /
 alkoxycarbonyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C>
- G3 = NH₂ / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G4) / CN / CO₂H /
 alkoxycarbonyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C>
- G4 = F / Cl / Br / I
- G5 = O / S / 16



- G6 = alkyl <containing 1-6 C> / OH / CN
- G7 = (up to 2) N / 18



- G8 = H / alkyl <containing 1-6 C> / NH₂ /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / F / Cl / Br / I
- G9 = O / S / NH / 20



- G10 = alkyl <containing 1-6 C>
- G11 = O / S / NH / 24 / 26

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR



EE 200300384	A	20031215	EE 2003-384	20020211
JP 2004523546	T2	20040805	JP 2002-564505	20020211
CN 1537105	A	20041013	CN 2002-805014	20020211
BR 2002007268	A	20050315	BR 2002-7268	20020211
US 2002193377	A1	20021219	US 2002-75954	20020213
ZA 2003006008	A	20041104	ZA 2003-6008	20030804
NO 2003003593	A	20030813	NO 2003-3593	20030813
BG 108091	A	20041230	BG 2003-108091	20030813

PRIORITY APPLN. INFO.:

US 2001-268661P	20010214
WO 2002-EP1979	20020211

OTHER SOURCE(S): CASREACT 137:185501


GI





AB Title compds. I [R1 = H, amino, alkyl, alkenyl, alkynyl, alkylamino, aryl, heterocycle, etc.; W = O, S, =N-R'; R' = alkyl, OH, CN; X1-3 = N, C-R6; R6 = H, alkyl, amino, alkylamino, etc.; Y = O, S, NH, N-alkyl; Z = O, S, NR7; R7 = H, alkyl, aryl, aryl, heteroaryl, etc.; n = 1-8; Z1 = alkyl; A = (non)aromatic, 5- or 6-membered monocycle comprising from 0 to 4 heteroatoms selected from N, O, S, etc.; m = 0-7; R2 = alkyl, halo, CN, NO2, SCF3, CF3, OCF3, etc.; R3 = H, alkyl, alkenyl, alkynyl, etc.] were prepared Over 200 synthetic examples were provided. For instance, di-Me 4-aminoisophthalate was reacted with benzylisocyanate and heated to 95-100° overnight to give Me 3-benzyl-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carboxylate which was saponified (dioxaneaq, LiOH, reflux) to give the carboxylic acid. This intermediate was coupled with benzylamine to afford II. Selected examples of I had IC50 = 2.25 - 0.001 μM for MMP13 and IC50 > 100 μM for MMP1, MMP2, MMP3, MMP7, MMP9, MMP12 and MMP14; II had IC50 = 0.193 μM for MMP13. Compds. I are useful for the treatment of osteoarthritis and rheumatoid arthritis.

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1



G34 = H / 256 / 260



G35 = R <"drug or drug bearing moiety">

G36 = H / alkyl / cycloalkyl / Ph (substd.)

G37 = alkylene <containing 1-12 C>

Patent location: claim 1

Note: also incorporates claim 9

Note: substitution is restricted

L86 ANSWER 20 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 137:185501 MARPAT

TITLE: Preparation of quinazolines as specific inhibitors of type-13 matrix metalloprotease

INVENTOR(S): Adrianjara, Charles; Chantel-Barvian, Nicole; Gaudilliere, Bernard; Jacobelli, Henri; Ortwine, Daniel Fred; Patt, William Chester; Pham, Ly; Kostlan, Catherine Rose; Wilson, Michael William

PATENT ASSIGNEE(S): Warner-Lambert Company, USA

SOURCE: PCT Int. Appl., 264 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002064572	A1	20020822	WO 2002-EP1979	20020211
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW:				
GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2437122	AA	20020822	CA 2002-2437122	20020211
EP 1368324	A1	20031210	EP 2002-722137	20020211
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				


G25 = H / OH / 130




G26 = H / 135 / 138





G27 = 143 / H



G28 = H / 146




G29 = 161-7 162-160 / 167-7 165-160 / 170-7 171-160 /
176-7 174-160 / O / S / S(O) / SO₂ / 178 / 181 /
alkylene <containing 1-12 C> / 320-7 321-160 /
322-7 323-160




G30 = R <"leaving or linking group,
or diagnostic or therapeutic agent">

G31 = O / 168




G32 = bond / SO₂






G33 = 191 / 254





G20 = H / R / (Specifically claimed: Me)
 G21 = ethynyl / propargyl
 G22 = 100-3 98-96 / bond




G23 = H / R <"leaving or linking group,
 or diagnostic or therapeutic agent"> / 127 / 132 / 141 / SH /
 148 / 150 / 154 / 157 / alkyl <containing 1-12 C>
 (opt. substd. by OH) / alkoxy <containing 1-12 C> / 159 /
 (Examples: 183 / 212 / 222 / 224 / 227 / 279)




G24 = CH / N




G12 = O / S
 G13 = CH2 (opt. substd.) / C(O)
 G14 = bond / O / 65




G15 = 70-68 72-7 / bond




G16 = bond / 73




G17 = bond / 79-76 81-7 / SO2




G18 = bond / 107-3 105-102



G19 = S / S(O) / SO2





G6 = O / S / 57



G7 = H / F / Cl / Br / I / At /
 alkyl <containing 1-12 C> / alkoxy <containing 1-12 C> /
 CHO / alkylcarbonyl <containing 1-11 C> /
 alkenyl <containing 2-12 C> / alkynyl <containing 2-12 C> /
 alkoxy carbonyl <containing 1-12 C> /
 alkylaminocarbonyl <containing 1-12 C> /
 dialkylaminocarbonyl <each alkyl containing 1-12 C> /
 (Specifically claimed: Me)






G8 = any ring <containing 0-3 heteroatoms,
 0-3 N (no other heteroatoms), 3 or more C,
 1 or more double bonds, attached through 3 or more C,
 6-membered monocyclic ring> (opt. substd. by 1 or more G9) /
 59 / (Specifically claimed: 122-2 121-4 124-6)



G9 = F / Cl / Br / I / At / alkyl <containing 1-12 C> /
 alkoxy <containing 1-12 C> / CHO /
 alkylcarbonyl <containing 1-11 C> /
 alkenyl <containing 2-12 C> / alkynyl <containing 2-12 C> /
 alkoxy carbonyl <containing 1-12 C> /
 alkylaminocarbonyl <containing 1-12 C> /
 dialkylaminocarbonyl <each alkyl containing 1-12 C>

G10 = any ring <containing 0-2 heteroatoms,
 0-2 N (no other heteroatoms), 4 or more C,
 1 or more double bonds, attached through 4 or more C,
 6-membered monocyclic ring> (opt. substd.)


G11 = CH2 (opt. substd.) / C(O) / 61-3 63-7 /
 67-3 69-7 / 75-3 77-7 / 82-3 83-7 / 84-3 85-7 /
 88-3 89-7 / 91-3 93-7 / 95-3 97-7 / 101-3 103-7 /
 108-3 109-7 / 111-3 110-7 / 114-3 113-7 / 117-3 119-7 /
 304-3 305-7 / 306-3 307-7 / 308-3 310-7 / 311-3 313-7 /
 314-3 316-7 / 317-3 319-7





DMSO, followed by addition of DTPA tris(tert-butyl) ester. The folate mimetic is conjugated to a diagnostic or therapeutic agent to enable selective delivery of the agent to the targeted cell population. The ^{111}In complex of II was selectively localized in the folate-receptor-pos. tumor xenografts of human KB cells in athymic mice (NuNu strain) and afforded prolonged tumor retention of ^{111}In (5.4, 5.5, and 3.6% ID/g at 1 h, 4 h, and 24 h, resp.); blockable binding was also observed in the kidneys, where the folate receptor occurs in the proximal tubes.

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


MSTR 1



G1 = phenylene (opt. substd. by 1 or more G2) /
 10-6 9-8 / 15-6 18-8 / 20-6 22-8 / 24-6 25-8 /
 29-6 33-8 / 34-6 37-8




G2 = F / Cl / Br / I / At / alkyl <containing 1-12 C> /
 alkoxy <containing 1-12 C> / (Specifically claimed: Me)
 G3 = S / O / 51




G4 = H / F / Cl / Br / I / At /
 alkyl <containing 1-12 C> / alkoxy <containing 1-12 C> /
 (Specifically claimed: Me)
 G5 = F / Cl / Br / I / At / H /
 alkyl <containing 1-12 C> / alkoxy <containing 1-12 C> /
 CHO / alkylcarbonyl <containing 1-11 C> /
 alkenyl <containing 2-12 C> / alkynyl <containing 2-12 C> /
 alkoxy carbonyl <containing 1-12 C> /
 alkylaminocarbonyl <containing 1-12 C> /
 dialkylaminocarbonyl <each alkyl containing 1-12 C> / 55 /
 (Specifically claimed: Me)

LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
 UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 EP 1389209 A1 20040218 EP 2002-731495 20020424
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
 US 2004242582 A1 20041202 US 2004-475876 20040621
 US 2005227985 A9 20051013
 PRIORITY APPLN. INFO.: US 2001-286082P 20010424
 WO 2002-US13045 20020424

GI




I




II

AB A cell population expressing folate receptors is selectively targeted with folate mimetic I [D = (CR6R7)s; E = (A1)p; F = (CR6R7)1-s; G = (A2)r; X, Y = halo, R2, OR2, SR3, NR4R5; U, V, W = (R6')C:, N:, (R6')C(R7'), N(R4')]; T = S, O, N, C:C (such that T is part of an aromatic ring); A1, A2 = C(Z), C(Z)O, OC(Z), N(R4'), C(Z)N(R4'), N(R4')C(Z), OC(Z)N(R4'), N(R4')C(Z)O, N(R4')C(Z)N(R5'), O, S, S(O), SO2, N(R4')SO2, C(R6')C(R7'), N(C.tplbond.CH), N(CH2C.tplbond.CH), C1-12-alkyl, C1-12-alkoxy; Z = O, S; R1 = H, halo, C1-12-alkyl, C1-12-alkoxy; R2 - R5, R4', R4'' - R7'' = H, halo, C1-12-alkyl, C1-12-alkoxy, C1-12-alkenyl, C1-12-alkynyl, (C1-12-alkoxy)carbonyl, (C1-12-alkylamino)carbonyl; R6, R7 = H, halo, C1-12-alkyl, C1-12-alkoxy; R6R7 = O; R6', R7' = H, halo, C1-12-alkyl, C1-12-alkoxy; R6'R7' = O; L = divalent linker (with the proviso that L ≠ naturally occurring amino acid covalently linked to A2 at its α-amino group through an amide bond); n, p, r, s = 0, 1; B = H, leaving group]. Thus, CYK4-013 (II) was prepared from pterioic acid via coupling with (CH2OCH2CH2NH2)2 using PyBOP, HOBT and N-methylmorpholine in






G35 = N / 118



G36 = H / R / (Specifically claimed: Me)

G37 = carbocycle <containing 5-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
(Specifically claimed: Ph / 143 / 150 / 157)



G38 = (0-3) CH₂

G39 = OH / alkoxy <containing 1-6 C> /
(Specifically claimed: OMe)

Patent location: claim 21

L86 ANSWER 19 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 137:338133 MARPAT

TITLE: Preparation of folate mimetics and folate-receptor
binding conjugates thereof

INVENTOR(S): Green, Mark A.; Leamon, Christopher P.; Ke, Chun-Yen

PATENT ASSIGNEE(S): Purdue Research Foundation, USA; Endocyte, Inc.

SOURCE: PCT Int. Appl., 52 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002085908	A1	20021031	WO 2002-US13045	20020424
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				



G25 = O / S / NH / 66



G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH2 / O / S / S(O) / SO2
 G28 = carbocycle <containing 5-6 C;
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29)
 G29 = alkyl <containing 1-6 C> / halo /
 alkyl <containing 1-6 C> (substd. by (3) halo) / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G30 = aryl <containing up to 10 C>
 (opt. substd. by 1 or more G4)
 G31 = cycloalkyl <containing 3-10 C>
 (opt. substd. by 1 or more G4)
 G32 = H / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G33) /
 alkenyl <containing 3-6 C> (opt. substd. by 1 or more G33) /
 alkynyl <containing 3-6 C> (opt. substd. by 1 or more G33) /
 97 / carbocycle <containing 5-10 C, 0 or more double bonds,
 mono- or bicyclic, 5- or 6-membered rings only>
 (opt. substd. by (1-7) G21) / heterocycle <containing 5-10
 atoms, 1-4 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21)






G33 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
 CO2H / alkoxycarbonyl <containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 99 / 101-20 103-98 /
 (Specifically claimed: CH2)

- G15 = halo / NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / OH / SH /
alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
CO2H / alkoxycarbonyl <containing 1-6 C>
G16 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
G18 = O / S / S(O) / SO2 / NH / 41

$\text{N} \text{---} \text{G19}$
41

- G19 = alkyl <containing 1-6 C>
G21 = alkyl <containing 1-6 C> / halo / CN / NO2 /
alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 43 /
45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
zero or more O (no other heteroatoms),
attached through 1 or more N, 5- to 6-membered monocyclic
ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
73 / 76 / CO2H / alkoxycarbonyl <containing 1-6 C> / 80 /
83 / 86 / 90 / 92 / 95 / carbocycle <containing 5-6 C,
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G29) /
heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, 5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G29)



- G22 = alkyl <containing 1-6 C> /
aryl <containing up to 10 C>
G23 = (1-3) CH2
G24 = NH2 / 56 / 58 / heterocycle <containing 1-2
heteroatoms, 1 or more N, zero or more O (no other
heteroatoms), attached through 1 or more N,
5- to 6-membered monocyclic ring>

zero or more O, zero or more S (no other heteroatoms),
monocyclic> (opt. substd. by 1 or more G4) /
heterocycle <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G4) / (Specifically claimed: Me)
G4 = halo / NH₂ / alkylamino <containing 1-10 C> /
dialkylamino <each alkyl containing 1-10 C> / CN /
alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
SH / alkoxy <containing 1-6 C> /
alkylthio <containing 1-6 C>

$\text{C}(\text{O})\text{G5}$
16

G5 = alkyl <containing 1-6 C> /
aryl <containing up to 10 C> / OH /
alkoxy <containing 1-16 C>
G6 = O / S / **NH** / 13

$\text{N}\text{---}\text{G2}$
13

G7 = (up to 2) N / 22

$\text{C}\text{---}\text{G8}$
22

G8 = H / alkyl <containing 1-6 C> / OH /
alkoxy <containing 1-6 C> / **halo** / CF₃ / CN / NO₂ / SH / 24
/
26 / NH₂ / 28 / 30 / heterocycle <containing 1-2
heteroatoms, 1 or more N, zero or more O (no other
heteroatoms), attached through 1 or more N,
5- to 6-membered monocyclic ring>

$\text{G9}\text{---}\text{OH}$
24

$\text{G10}\text{---}\text{G11}$
26

$\text{HN}\text{---}\text{G12}$
28




$\text{N}\begin{matrix} \text{G12} \\ \text{30} \\ \text{G12} \end{matrix}$

G9 = S / S(O)
G10 = S / S(O) / SO₂
G11 = alkyl <containing 1-6 C>
G12 = alkyl <containing 1-6 C> /
alkyl <containing 1-10 C> (substd. by 1 or more aryl
<containing up to 10 C>)
G13 = H / halo / OSO₂Me / 168 / CHO / COMe /
R <"ester group"> / ethynyl / 171 / (Specifically claimed: I)

$\text{O}\text{---}\text{SO}_2\text{---}\text{CF}_3$
168

$\begin{matrix} \text{Br} \\ | \\ \text{HC}=\text{C}\text{---}\text{Br} \\ 171 \end{matrix}$

5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 (Specifically claimed: Ph / 143 / 150 / 157)



G38 = (0-3) CH₂

G39 = OH / alkoxy <containing 1-6 C> /
 (Specifically claimed: OMe)

Patent location:

claim 1

Note:

and N-oxides or pharmaceutically acceptable acid or
 base addition salts


Note:

additional heteroatom interruptions in G17 also
 claimed



Stereochemistry:

and optical isomers

MSTR 2



G1 = 1-19 9-21 / 104-19 112-21



G2 = alkyl <containing 1-6 C> / OH / CN

G3 = H / CF₃ / NH₂ / alkylamino <containing 1-10 C> /
 dialkylamino <each alkyl containing 1-10 C> /
 alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
 alkenyl <containing 3-6 C> (opt. substd. by 1 or more G4) /
 alkynyl <containing 3-6 C> (opt. substd. by 1 or more G4) /
 aryl <containing up to 10 C> (opt. substd. by 1 or more G4) /
 alkyl <containing 1-10 C> (substd. by 1 or more G30) /
 alkyl <containing 1-10 C> (substd. by G31) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,

G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH2 / O / S / S(O) / SO2
 G28 = carbocycle <containing 5-6 C,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29)
 G29 = alkyl <containing 1-6 C> / halo /
 alkyl <containing 1-6 C> (substd. by (3) halo) / OH /
 alkoxy <containing 1-6 C> / SH /
 alkylthio <containing 1-6 C> / NH2 /
 alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C>
 G30 = aryl <containing up to 10 C>
 (opt. substd. by 1 or more G4)
 G31 = cycloalkyl <containing 3-10 C>
 (opt. substd. by 1 or more G4)
 G32 = H / alkyl <containing 1-6 C>
 (opt. substd. by 1 or more G33) /
 alkenyl <containing 3-6 C> (opt. substd. by 1 or more G33) /
 alkynyl <containing 3-6 C> (opt. substd. by 1 or more G33) /
 97 / carbocycle <containing 5-10 C, 0 or more double bonds,
 mono- or bicyclic, 5- or 6-membered rings only>
 (opt. substd. by (1-7) G21) / heterocycle <containing 5-10
 atoms, 1-4 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21)

~~G34-G37~~
 97 98


G33 = NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
 CO2H / alkoxycarbonyl <containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
 G34 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 99 / 101-20 103-98 /
 (Specifically claimed: CH2)

~~G16=O~~ ~~G17-G18-G17~~
 99 101 103




G35 = N / 118

~~C-G36~~
 118

G36 = H / R / (Specifically claimed: Me)
 G37 = carbocycle <containing 5-10 C,
 0 or more double bonds, mono- or bicyclic,




G21 = alkyl <containing 1-6 C> / halo / CN / NO2 /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 43 /
 45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
 73 / 76 / CO2H / alkoxy carbonyl <containing 1-6 C> / 80 /
 83 / 86 / 90 / 92 / 95 / carbocycle <containing 5-6 C,
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29) /
 heterocycle <containing 5-6 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G29)




G22 = alkyl <containing 1-6 C> /
 aryl <containing up to 10 C>

G23 = (1-3) CH2


G24 = NH2 / 56 / 58 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>




G25 = O / S / NH / 66




alkoxy <containing 1-6 C> / **halo** / CF₃ / CN / NO₂ / SH / 24
 /
 26 / NH₂ / 28 / 30 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>




G9 = S / S(O)
 G10 = S / S(O) / SO₂
 G11 = alkyl <containing 1-6 C>
 G12 = alkyl <containing 1-6 C> /
 alkyl <containing 1-10 C> (substd. by 1 or more aryl
 <containing up to 10 C>)
 G13 = ethynylene / 33-18 35-20




G14 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd. by 1 or more G15) / 36 / 38-18 40-34 /
 (Specifically claimed: CH₂)




G15 = halo / NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
 CO₂H / alkoxycarbonyl <containing 1-6 C>
 G16 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G17 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)
 G18 = O / S / S(O) / SO₂ / NH / 41




G19 = alkyl <containing 1-6 C>
 G20 = carbocycle <containing 5-10 C,
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
 zero or more N, zero or more O,
 zero or more S (no other heteroatoms),
 0 or more double bonds, mono- or bicyclic,
 5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
 (Specifically claimed: Ph / 122 / 129 / 136)




G1 = 1-19 9-21 / 104-19 112-21




G2 = alkyl <containing 1-6 C> / OH / CN
 G3 = H / CF3 / NH2 / alkylamino <containing 1-10 C> /
 dialkylamino <each alkyl containing 1-10 C> /
 alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
 alkenyl <containing 3-6 C> (opt. substd. by 1 or more G4) /
 alkynyl <containing 3-6 C> (opt. substd. by 1 or more G4) /
 aryl <containing up to 10 C> (opt. substd. by 1 or more G4) /
 alkyl <containing 1-10 C> (substd. by 1 or more G30) /
 alkyl <containing 1-10 C> (substd. by G31) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 monocyclic> (opt. substd. by 1 or more G4) /
 heterocycle <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G4) / (Specifically claimed: Me)
 G4 = halo / NH2 / alkylamino <containing 1-10 C> /
 dialkylamino <each alkyl containing 1-10 C> / CN /
 alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
 SH / alkoxy <containing 1-6 C> /
 alkylthio <containing 1-6 C>





G5 = alkyl <containing 1-6 C> /
 aryl <containing up to 10 C> / OH /
 alkoxy <containing 1-16 C>
 G6 = O / S / NH / 13



G7 = (up to 2) N / 22



G8 = H / alkyl <containing 1-6 C> / OH /



AB Title compds. I [wherein A = (hetero)aryl or (hetero)cycloalkyl; W1 = O, S, or NR₃; W2 = H, CF₃, NH₂, (di)alkylamino, or (un)substituted (cycloalkyl)alkyl, alkenyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; or W1W2 = NX₄=W₃; W₃ = N or CR₅; X₁-X₃ = independently N or (un)substituted C; X₄ = N or CR₇; X₅ = O, S, NH, or N-alkyl; X₆ = bond, CH₂, O, SO₀₋₂; Z = CR₁₂R₁₃; R₁ = H, alkyl, alkenyl, alkynyl, or (un)substituted (hetero)aryl or (hetero)cycloalkyl; R₂ = independently H, (trihalo)alkyl, halo, CN, NO₂, (CH₂)_kNR₁₀R₁₁, OR₁₄, SR₁₄, SO₂R₁₄, acyl, X₅(CH₂)_kNR₁₀R₁₁, X₅(CH₂)_kSO₂NR₁₄R₁₅, X₅(CH₂)_kCO₂R₁₄, (CH₂)_kCO₂R₁₄, X₅(CH₂)_kCONR₁₄R₁₅, (CH₂)_kCONR₁₄R₁₅ X₆R₁₆, and trialkylsiloxy; R₃ = H, alkyl, OH, or CN; R₄ = H or alkyl; R₅ = H, OR₆, SR₆, or (un)substituted (cyclo)alkyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; R₆, R₈, and R₉ = independently H or (aryl)alkyl; R₇ = H, NR₈R₉, OR₈, SR₈, or (un)substituted (cyclo)alkyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; R₁₀ and R₁₁ = independently H, (hydroxy)alkyl, or arylalkyl; or NR₁₀R₁₁ = (un)substituted heterocyclalkyl; R₁₂ and R₁₃ = independently H, (trihalo)alkyl, halo, NH₂, (di)alkylamino, OR₄, SR₄, or CO₂R₄; R₁₄ and R₁₅ = independently H or alkyl; R₁₆ = (un)substituted (hetero)aryl or (hetero)cycloalkyl; k = 0-3; n = 0-8; q = 0-7; with provisos; or isomers, N-oxides, or pharmaceutically acceptable salts thereof] were prepared as specific inhibitors of type 13 matrix metalloprotease (MMP-13). For example, reaction of Me 4-(aminomethyl)benzoate•HCl with 2-amino-5-iodobenzoic acid using DEC•HCl and TEA in DMF provided the amide (70%). Cyclization using 1,1'-carbonyldiimidazole in THF gave the quinazoline (99.5%), which was methylated using MeI in the presence of K₂CO₃ in DMF to afford Me 4-(6-iodo-1-methyl-2,4,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)methylbenzoate (64.2%). Substitution with 3-(4-methoxyphenyl)prop-1-yne catalyzed by Pd(PPh₃)₂Cl₂ and CuI in TEA gave II (6%). Invention compds. inhibited the proteolysis of a peptide substrate with MMP-13 with IC₅₀ values <1 μM, generally 100 times lower than the IC₅₀ values for the same compds. with respect to MMP-1, MMP-2, MMP-3, MMP-7, MMP-9, MMP-12, and MMP-14. Thus, I are useful for the treatment of arthritis, cancer, and other diseases mediated by MMP-13 (no data).

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1

INVENTOR(S): Gaudilliere, Bernard; Jacobelli, Henry
 PATENT ASSIGNEE(S): Warner-Lambert Company, USA
 SOURCE: PCT Int. Appl., 68 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003033477	A1	20030424	WO 2001-EP11824	20011012
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2463159	AA	20030424	CA 2002-2463159	20021011
WO 2003033478	A1	20030424	WO 2002-EP12194	20021011
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2003130278	A1	20030710	US 2002-269197	20021011
US 6962922	B2	20051108		
BR 2002013239	A	20040928	BR 2002-13239	20021011
EP 1465878	A1	20041013	EP 2002-801341	20021011
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK			
JP 2005509626	T2	20050414	JP 2003-536218	20021011
US 2005245548	A1	20051103	US 2005-148880	20050609
PRIORITY APPLN. INFO.:			US 2001-329181P	20011012
			WO 2001-EP11824	20011012
			US 2002-395441P	20020712
			WO 2002-EP8475	20020712
			US 2002-269197	20021011
			WO 2002-EP12194	20021011

GI

G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)
G18 = O / S / S(O) / SO2 / NH / 41

$\text{N} \text{---} \text{G19}$
41

G19 = alkyl <containing 1-6 C>
G30 = aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4)
G31 = carbocycle <containing 3-10 C, non-aromatic,
0 or more double bonds, mono- or bicyclic>
(opt. substd. by 1 or more G4)
G32 = H / alkyl <containing 1-6 C> (opt. substd.) /
alkenyl <containing 2-6 C> (opt. substd.) /
alkynyl <containing 2-6 C> (opt. substd.) / 97 /
carbocycle <containing 5-10 C, 0 or more double bonds,
mono- or bicyclic, 5- or 6-membered rings only>
(opt. substd.) / heterocycle <containing 5-10 atoms,
1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd.)

$\text{G34} \text{---} \text{G37}$
97 98

G34 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by 1 or more G15) / 99 / 101-20 103-98

$\text{G16} \text{=O}$ $\text{G17} \text{---} \text{G18} \text{---} \text{G17}$
99 101 103

G35 = N / 118

$\text{C} \text{---} \text{G36}$
118

G36 = H / R
G37 = carbocycle <containing 5-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd.) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd.)

Patent location: claim 24

L86 ANSWER 18 OF 26 MARPAT COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 138:338159 MARPAT

TITLE: Preparation of alkynylated fused ring pyrimidine
compounds as matrix metalloprotease 13 inhibitors

dialkylamino <each alkyl containing 1-6 C> / CN /
 alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /
 SH / alkoxy <containing 1-6 C> /
 alkylthio <containing 1-6 C>

$\overset{\text{C}(\text{O})\text{-G5}}{\underset{16}{\text{C}}}$

G5 = alkyl <containing 1-6 C> / Ph / OH /
 alkoxy <containing 1-16 C>
 G6 = O / S / **NH** / 13

$\overset{\text{N}-\text{G2}}{\underset{13}{\text{N}}}$

G7 = (up to 2) N / 22

$\overset{\text{C}-\text{G8}}{\underset{22}{\text{C}}}$

G8 = H / alkyl <containing 1-6 C> / OH /
 alkoxy <containing 1-6 C> / **halo** / CF3 / CN / NO2 / SH / 24
 /
 26 / NH2 / 28 / 30 / heterocycle <containing 1-2
 heteroatoms, 1 or more N, zero or more O (no other
 heteroatoms), attached through 1 or more N,
 5- to 6-membered monocyclic ring>

$\overset{\text{G9}-\text{OH}}{\underset{24}{\text{G9}}}$



$\overset{\text{G10}-\text{G11}}{\underset{26}{\text{G10}}}$

$\overset{\text{HN}-\text{G12}}{\underset{28}{\text{HN}}}$


$\begin{array}{c} \text{G12} \\ \diagup \\ \text{N} \\ \diagdown \\ \text{G12} \end{array}$
 30

G9 = S / S(O)
 G10 = S / S(O) / SO2
 G11 = alkyl <containing 1-6 C>
 G12 = alkyl <containing 1-6 C> /
 alkyl <containing 1-6 C> (substd. by 1 or more aryl
 <containing up to 10 C, mono- or bicyclic>)
 G13 = H / halo / OSO2Me / 344 / CHO / COMe /
 R <"ester group"> / 347 / ethynyl / 353

$\overset{\text{O}-\text{SO}_2\text{-CF}_3}{\underset{344}{\text{O}}}$



G15 = halo / NH2 / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> / OH / SH /
 alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C> /
 CO2H / alkoxy carbonyl <containing 1-6 C>
 G16 = carbon chain <containing 1 or more C,
 0 or more double bonds, 0 or more triple bonds>
 (opt. substd.)



G50 = Cl / F

Patent location:

Note:

Note:

Note:

Stereochemistry:


claim 1

and N-oxides or pharmaceutically acceptable acid or base addition salts


additional heteroatom interruptions in G17 also claimed

substitution is restricted

and optical isomers

MSTR 2


G1 = 1-19 9-21 / 104-19 112-21




G2 = alkyl <containing 1-6 C> / OH / CN

G3 = H / CF₃ / NH₂ / alkylamino <containing 1-6 C> /
 dialkylamino <each alkyl containing 1-6 C> /
 alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
 alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
 alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
 aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by 1 or more G4) /
 alkyl <containing 1-6 C> (substd. by 1 or more G30) /
 alkyl <containing 1-6 C> (substd. by G31) /
 heteroaryl <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 monocyclic> (opt. substd. by 1 or more G4) /
 heterocycle <containing 1-4 heteroatoms, zero or more N,
 zero or more O, zero or more S (no other heteroatoms),
 non-aromatic, 0 or more double bonds,
 5- to 6-membered monocyclic ring>
 (opt. substd. by 1 or more G4)


G4 = halo / NH₂ / alkylamino <containing 1-6 C> /




G42 = alkylene <containing 1-6 C>
 G43 = OH / alkoxy <containing 1-6 C> / NH2 / 220 / 222 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring> / 225




G44 = OH / alkoxy <containing 1-6 C> / NH2 / 227 / 229 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring>




G45 = 237 / OH / alkoxy <containing 1-6 C> / 240 / (Specifically claimed: CO2Me)




G46 = alkylene <containing 2-4 C>
 G47 = NH2 / 244 / 246 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring>







G48 = Ph / F / 263 / Cl



G49 = F / Cl / Br / CO2Bu-t / 291 / 298 / 302 / 309 / CH2OH / 313 / 321



G38 = (0-3) CH2
 G40 = alkyl <containing 1-6 C>
 G41 = alkyl <containing 1-6 C> / halo / CN / NO2 /
 alkyl <containing 1-6 C> (substd. by (3) halo) / NH2 / 232 /
 234 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
 zero or more O (no other heteroatoms),
 attached through 1 or more N, 5- to 6-membered monocyclic
 ring> / OH / SH / 173 / 175 / 177 / 179 / 181 / 183 / 186 /
 189 / 191 / 194 / CO2H / alkoxy carbonyl <containing 1-6 C> /
 198 / 201 / 204 / 208 / 210 / 213 / 215 /
 aryl <containing up to 10 C, mono- or bicyclic>
 (opt. substd. by (1-4) G29) / heteroaryl <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), mono- or bicyclic>
 (opt. substd. by (1-4) G29) / heterocycle <containing 1-4
 heteroatoms, zero or more N, zero or more O,
 zero or more S (no other heteroatoms), non-aromatic,
 0 or more double bonds, mono- or bicyclic,
 (1) 5- or more membered ring, (1) up to 6-membered ring>
 (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,
 non-aromatic, 0 or more double bonds, mono- or bicyclic>
 (opt. substd. by 1 or more G29)



- non-aromatic, 0 or more double bonds, mono- or bicyclic>
(opt. substd. by 1 or more G29)
- G29 = alkyl <containing 1-6 C> / halo /
alkyl <containing 1-6 C> (substd. by (3) halo) / OH /
alkoxy <containing 1-6 C> / SH /
alkylthio <containing 1-6 C> / NH2 /
alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C>
- G30 = aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4)
- G31 = carbocycle <containing 3-10 C, non-aromatic,
0 or more double bonds, mono- or bicyclic>
(opt. substd. by 1 or more G4)
- G32 = H / alkyl <containing 1-6 C>
(opt. substd. by 1 or more G33) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G33) /
97 / carbocycle <containing 5-10 C, 0 or more double bonds,
mono- or bicyclic, 5- or 6-membered rings only>
(opt. substd. by (1-7) G41) / heterocycle <containing 5-10
atoms, 1-4 heteroatoms, zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41)

~~G34-G37~~
97 98





- G33 = NH2 / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> /
alkyl <containing 1-6 C> (opt. substd. by (3) halo) / CN /
CO2H / alkoxy carbonyl <containing 1-6 C> / OH / SH /
alkoxy <containing 1-6 C> / alkylthio <containing 1-6 C>
- G34 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd. by 1 or more G15) / 99 / 101-20 103-98 /
(Specifically claimed: CH2)

~~G16=O~~ ~~G17-G18-G17~~
99 101 103


- G35 = N / 118

~~C~~-G36
118


- G36 = H / R / (Specifically claimed: Me)
- G37 = carbocycle <containing 5-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
heterocycle <containing 5-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G41) /
(Specifically claimed: Ph / 143 / 150 / 157 / 251 / 331 /
337)



G22 = alkyl <containing 1-6 C> / Ph
 G23 = (1-3) CH₂
 G24 = NH₂ / 56 / 58 / heterocycle <containing 1-2 heteroatoms, 1 or more N, zero or more O (no other heteroatoms), attached through 1 or more N, 5- to 6-membered monocyclic ring>



G25 = O / S / NH / 66




G26 = OH / alkoxy <containing 1-6 C>
 G27 = CH₂ / O / S / S(O) / SO₂
 G28 = aryl <containing up to 10 C, mono- or bicyclic> (opt. substd. by (1-4) G29) / heteroaryl <containing 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), mono- or bicyclic> (opt. substd. by (1-4) G29) / heterocycle <containing 1-4 heteroatoms, zero or more N, zero or more O, zero or more S (no other heteroatoms), non-aromatic, 0 or more double bonds, mono- or bicyclic, (1) 5- or more membered ring, (1) up to 6-membered ring> (opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,

0 or more double bonds, 0 or more triple bonds>
(opt. substd.)




G17 = carbon chain <containing 1 or more C,
0 or more double bonds, 0 or more triple bonds>
(opt. substd.)

G18 = O / S / S(O) / SO₂ / NH / 41

N—G19
41

G19 = alkyl <containing 1-6 C>

G20 = carbocycle <containing 8-10 C,
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
heterocycle <containing 8-10 atoms, 1-4 heteroatoms,
zero or more N, zero or more O,
zero or more S (no other heteroatoms),
0 or more double bonds, mono- or bicyclic,
5- or 6-membered rings only> (opt. substd. by (1-7) G21) /
(Specifically claimed: Ph / 122 / 258 / 270 / 275 / 280 /
285)



G21 = alkyl <containing 1-6 C> / halo / CN / NO₂ /
alkyl <containing 1-6 C> (substd. by (3) halo) / NH₂ / 43 /
45 / heterocycle <containing 1-2 heteroatoms, 1 or more N,
zero or more O (no other heteroatoms),
attached through 1 or more N, 5- to 6-membered monocyclic
ring> / OH / SH / 48 / 50 / 52 / 54 / 61 / 63 / 68 / 71 /
73 / 76 / CO₂H / alkoxy carbonyl <containing 1-6 C> / 80 /
83 / 86 / 90 / 92 / 95 / aryl <containing up to 10 C,
mono- or bicyclic> (opt. substd. by (1-4) G29) /
heteroaryl <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
mono- or bicyclic> (opt. substd. by (1-4) G29) /
heterocycle <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds, mono- or bicyclic,
(1) 5- or more membered ring, (1) up to 6-membered ring>
(opt. substd. by (1-4) G29) / carbocycle <containing 3-10 C,
non-aromatic, 0 or more double bonds, mono- or bicyclic>
(opt. substd. by 1 or more G29) / 168

¹⁶C (O)-G5

13 N—G2

C—G8
22

$$\begin{array}{ccc} \text{G}^9\text{---OH} & \text{G}^{10}\text{---G}^{11} & \text{HN---G}^{12} \\ 24 & 26 & 28 \end{array} \quad \begin{array}{c} \text{G}^{12} \\ \diagup \\ \text{N} \\ \diagdown \\ \text{G}^{12} \\ 30 \end{array}$$
$$\begin{array}{ccccc} & \text{G14} & - & \text{C} & \equiv & \text{C} \\ & \text{33} & & \text{34} & & \text{35} \end{array}$$
$$\begin{array}{ccc} \text{G16}=\text{O} & & \text{G17}-\text{G18}-\text{G17} \\ 36 & & 38 \qquad \qquad 40 \end{array}$$

Page 122



(trihalo)alkyl, halo, NH₂, (di)alkylamino, OR₄, SR₄, or CO₂R₄; R₁₄ and R₁₅ = independently H or alkyl; R₁₆ = (un)substituted (hetero)aryl or (hetero)cycloalkyl; k = 0-3; n = 0-8; q = 0-7; with provisos; or isomers, N-oxides, or pharmaceutically acceptable salts thereof] were prepared as specific inhibitors of type 13 matrix metalloprotease (MMP-13). For example, reaction of Me 4-(aminomethyl)benzoate•HCl with 2-amino-5-iodobenzoic acid using DEC•HCl and TEA in DMF provided the amide (70%). Cyclization using 1,1'-carbonyldiimidazole in THF gave the quinazoline (99.5%), which was methylated using MeI in the presence of K₂CO₃ in DMF to afford Me 4-(6-iodo-1-methyl-2,4,4-dioxo-1,4-dihydro-2H-quinazolin-3-ylmethyl)benzoate (64.2%). Substitution with 3-(4-methoxyphenyl)prop-1-yne catalyzed by Pd(PPh₃)₂Cl₂ and CuI in TEA gave II (6%). Invention compds. inhibited the proteolysis of a peptide substrate with MMP-13 with IC₅₀ values <10 μM, generally 100 times lower than the IC₅₀ values for the same compds. with respect to MMP-1, MMP-2, MMP-3, MMP-7, MMP-9, MMP-12, and MMP-14. Thus, I are useful for the treatment of arthritis, cancer, and other diseases mediated by MMP-13 (no data).

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1

G20-G13-G1-G32
18 19 20 21



G1 = 1-19 9-21 / 104-19 112-21





G2 = alkyl <containing 1-6 C> / OH / CN
G3 = H / CF₃ / NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> /
alkyl <containing 1-6 C> (opt. substd. by 1 or more G4) /
alkenyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
alkynyl <containing 2-6 C> (opt. substd. by 1 or more G4) /
aryl <containing up to 10 C, mono- or bicyclic>
(opt. substd. by 1 or more G4) /
alkyl <containing 1-6 C> (substd. by 1 or more G30) /
alkyl <containing 1-6 C> (substd. by G31) /
heteroaryl <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
monocyclic> (opt. substd. by 1 or more G4) /
heterocycle <containing 1-4 heteroatoms, zero or more N,
zero or more O, zero or more S (no other heteroatoms),
non-aromatic, 0 or more double bonds,
5- to 6-membered monocyclic ring>
(opt. substd. by 1 or more G4) / (Specifically claimed: Me)
G4 = halo / NH₂ / alkylamino <containing 1-6 C> /
dialkylamino <each alkyl containing 1-6 C> / CN /
alkyl <containing 1-6 C> (substd. by (3) halo) / 16 / OH /

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, US, UZ, VN, YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,
 FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF,
 CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 CA 2463159 AA 20030424 CA 2002-2463159 20021011
 BR 2002013239 A 20040928 BR 2002-13239 20021011
 EP 1465878 A1 20041013 EP 2002-801341 20021011
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK
 JP 2005509626 T2 20050414 JP 2003-536218 20021011
 PRIORITY APPLN. INFO.: WO 2001-EP11824 20011012
 WO 2002-EP8475 20020712
 WO 2002-EP12194 20021011

GI



AB Title compds. I [wherein A = (hetero)aryl or (hetero)cycloalkyl; W1 = O, S, or NR3; W2 = H, CF3, NH2, (di)alkylamino, or (un)substituted (cycloalkyl)alkyl, alkenyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; or W1W2 = NX4=W3; W3 = N or CR5; X1-X3 = independently N or (un)substituted C; X4 = N or CR7; X5 = O, S, NH, or N-alkyl; X6 = bond, CH2, O, or SO0-2; Z = CR12R13; R1 = H, alkyl, alkenyl, alkynyl, or (un)substituted (hetero)aryl or (hetero)cycloalkyl; R2 = independently H, (trihalo)alkyl, halo, CN, NO2, (CH2)kNR10R11, OR14, SR14, SO2R14, acyl, X5(CH2)kNR10R11 (CH2)kSO2NR14R15, X5(CH2)kCO2R14, (CH2)kCO2R14, X5(CH2)kCONR14R15, (CH2)kCONR14R15 X6R16, and trialkylsiloxy; R3 = H, alkyl, OH, or CN; R4 = H or alkyl; R5 = H, OR6, SR6, or (un)substituted (cyclo)alkyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; R6, R8, and R9 = H or (aryl)alkyl; R7 = H, NR8R9, OR8, SR8, or (un)substituted (cyclo)alkyl, (hetero)aryl, arylalkyl, or heterocyclalkyl; R10 and R11 = independently H, (hydroxy)alkyl, or arylalkyl; or NR10R11 = (un)substituted heterocyclalkyl; R12 and R13 = independently H,




G9 = Ph (opt. substd. by 1 or more G10)
 G10 = F / Cl / Br / I / NO₂ /
 alkoxy <containing 1-6 C> (opt. substd. by 1 or more G11) /
 alkyl <containing 1-6 C> (opt. substd. by 1 or more G11) /
 Ph (opt. substd. by 1 or more alkyl <containing 1-6 C>)
 G11 = F / Cl / Br / I
 G12 = (1-3) CH₂
 G13 = O / S / NH
 G14 = H / F / Cl / Br / I / alkyl <containing 1-6 C> /
 alkoxy <containing 1-6 C>
 Patent location: claim 1
 Note: substitution is restricted
 Note: or prodrugs, or salts

L86 ANSWER 17 OF 26 MARPAT COPYRIGHT 2006 ACS on STN


ACCESSION NUMBER: 138:338166 MARPAT
 TITLE: Preparation of alkynylated fused ring pyrimidine
 compounds as matrix metalloprotease 13 inhibitors
 INVENTOR(S): Gaudilliere, Bernard; Jacobelli, Henry; Wilson,
 Michael William; Picard, Joseph Armand
 PATENT ASSIGNEE(S): Warner-Lambert Company LLC, USA
 SOURCE: PCT Int. Appl., 99 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003033478	A1	20030424	WO 2002-EP12194	20021011
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2003033477	A1	20030424	WO 2001-EP11824	20011012
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2004007469	A1	20040122	WO 2002-EP8475	20020712
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				


zero or more S (no other heteroatoms),
 attached through 2 or more C, 1 or more double bonds,
 5- to 7-membered monocyclic ring> (opt. substd.) /
 25-1 26-3 / 47-1 48-3 / 52-1 53-3 / 57-1 58-3 /
 (Specifically claimed: 31-1 32-3 / 36-1 37-3 / 42-1 43-3)



G2 = F / Cl / Br / I / alkyl <containing 1-6 C> /
 alkoxy <containing 1-6 C>
 G3 = 9-4 10-2 10-7 / 11-4 12-2 11-7 / 13-4 14-2 14-7 /
 15-4 16-2 15-7



G4 = alkylene <containing 1-6 C> /
 alkenylene <containing 2-6 C> /
 cycloalkylene <containing 3-6 C> /
 cycloalkenylene <containing 3-6 C> /
 heterocycle <containing 1 or more N, zero or more O,
 zero or more S, 0 or more double bonds, monocyclic> /
 17-3 18-8




G5 = NH / 20



G6 = alkyl <containing 1-6 C>
 G7 = alkylene <containing 1-6 C> /
 alkenylene <containing 2-6 C>
 G8 = heterocycle <containing 1 or more N,
 zero or more O, zero or more S, attached through 1 or more N>
 / carbocycle / NH / (Specifically claimed: 80-7 77-21 /
 86-7 83-21 / 92-7 89-21 / heterocycle <containing 6 atoms,
 1-2 heteroatoms, 1-2 N (no other heteroatoms), non-aromatic,
 0-1 double bond, 6-membered monocyclic ring>)

PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA,
 UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 CA 2474434 AA 20030807 CA 2003-2474434 20030127
 EP 1469854 A1 20041027 EP 2003-703053 20030127
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
 US 2005080096 A1 20050414 US 2003-501334 20030127
 JP 2005516053 T2 20050602 JP 2003-563564 20030127
 AU 2002-197 20020129
 WO 2003-JP708 20030127
 PRIORITY APPLN. INFO.:


GI



AB The title compds. [I; R1 = H, halo, alkyl or alkoxy; A and two adjacent carbon atoms of the six membered ring to be bonded with A form benzene ring, pyridine ring, etc.; Y1:Y2 = N:C(L11R21), C(L12R22):N, CH:C(L13R23), C(L14R24):CH (wherein L11, L12, L13, L14 = alkylene, alkenylene, etc.; R21, R22, R23 and R24 = cyclic amino group, carbocyclic group or amino group which are substituted with (un)substituted Ph); provided that when A and two adjacent carbon atoms of the six membered ring to be bonded with A form benzene ring, then Y1:Y2 = C(L12R22):N, CH:C(L13R23), C(L14R24):CH] having poly(adenosine 5'-diphospho-ribose)polymerase (PARP) inhibitory activity, were prepared Thus, reacting 4-(4-phenyl-3,6-dihydro-1(2H)-pyridyl)butanimidamide with cyclohexanone-2-carboxylic acid Et ester in the presence of K2CO3 in EtOH afforded 2-[3-(4-phenyl-3,6-dihydro-1(2H)-pyridyl)propyl]-5,6,7,8-tetrahydro-4(3H)-quinazolinone which showed IC50 of < 0.5 μ M against human PARP.


REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

MSTR 1



G1 = o-C6H4 (opt. substd. by G2) /
 carbocycle <containing 5-7 C, attached through 2 or more C,
 1 or more double bonds, 5- to 7-membered monocyclic ring>
 (opt. substd. by G2) / heterocycle <containing 5-7 atoms,
 zero or more N, zero or more O,

Molecular Weight (MW): 328.33
 Lawson Number (LN): 30121, 2817
 File Segment (FS): Stereo compound
 Compound Type (CTYPE): heterocyclic
 Constitution ID (CONSID): 527267
 Tautomer ID (TAUTID): 514815
 Beilstein Citation (BSO): 5-26
 Entry Date (DED): 1988/11/28
 Update Date (DUPD): 1992/01/31



Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	2
FS	File Segment	1
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
DED	Entry Date	1
DUPD	Update Date	1
MP	Melting Point	1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
RX	Reaction Documents	1

RXPRO Substance is Reaction Product 1

All References:

ALLREF

1. Lalezari; Sadeghi-Milani, J.Heterocycl.Chem., CODEN: JHTCAD, 16,
<1979>, 707,708

=>